



Statement of biodiversity priorities

Part two priorities and actions

Contents

Chapter five

Our vision for nature recovery.....04

What can you do to help
achieve the vision?.....05

Strategic aims for nature recovery.....06

Chapter six

Priority natural landscapes.....10

Landscape-scale nature recovery.....10

Identification of priority
natural landscapes.....11

Beyond the priority natural
landscapes.....15

Nature recovery on our farmland.....15

The Fens.....15

The Claylands.....16

The Chalklands.....17

Nature recovery in our cities
towns and villages.....18

Chapter seven

Priorities for nature recovery.....20

Habitats identified as local priorities.....20

Habitat actions.....21

Species identified as local priorities.....46

Species actions.....46

Chapter eight

Local habitat map.....82

Habitat priorities / Methodology.....82

Next Steps.....83

Acknowledgements.....84

Photography Acknowledgments.....85





5

Our vision for nature recovery



Our vision for nature recovery

Creating a wildlife-rich, resilient, productive and sustainable landscape, for people and nature.

At the heart of Cambridgeshire and Peterborough's vision for nature recovery is taking action to create, restore, connect and ultimately double the amount of nature. This will enhance our rich natural and agricultural heritage.

Doubling nature will see internationally significant wildlife sites extended, buffered, enriched and connected through nature-filled corridors and stepping stones across our landscape. This means that land managed for nature will increase from around 8% today to more than 16% by 2040.

The ambition to at least double nature in the area is viewed as an achievable target. This strategy goes further, identifying just under 30% of land as providing the best opportunities for nature restoration activity. This provides scope to exceed our doubling nature target beyond 2040.

The vision sees the creation of corridors of abundant biodiversity threading between and through our villages, towns and cities. It envisages vibrant communities taking actions for nature in gardens, parks and local wild spaces.

As an area with a strong growth agenda, the vision is for new high-quality nature-rich areas to be integrated with housing and commercial developments. Newly created natural green spaces within easy reach of communities will help attract businesses and retain staff to benefit our local economy.

In the fens and river valleys, functioning wetlands will connect across the landscape through wildlife-rich rivers, ditches, drains and floodplains. Improved water management will support our internationally important habitats and our productive agricultural land. Nature recovery will take place alongside securing water supplies for our existing and growing population, while also providing flood relief in affected areas. Identifying and effectively managing restorable peatlands will also increase the longevity and resilience of productive land and reduce carbon emissions.



Our vision for nature recovery (continued)

Our chalk springs and streams will be restored to provide diverse habitats for some of our most precious species as well as creating tranquil places for people to connect with nature. Our historic orchards will be valued, and new community orchards established to provide fruit for local communities. This will strengthen the connection with the area's fruit growing and horticultural heritage.

The farmed landscape of Cambridgeshire and Peterborough supports nationally important populations of farmland birds, insects and rare plants. Maintaining flower-rich field margins and increasing habitat connections through the sensitive management of ditches and

hedgerows, will create a diversity of wildlife-rich areas within a productive arable system.

In one of the driest areas of the country, our distinctive grasslands, woodlands, and wetland habitats will be cherished and managed to support the area's resilience to future changes in climate. Declines in species will be halted, and thriving, reconnected populations of plants and animals will be established.


Working alongside new approaches in sustainable development and farming, there will be more spaces for nature which will be made better, bigger and joined up, to ensure Cambridgeshire and Peterborough becomes an exemplar of nature recovery.

What can you do to help achieve the vision?

- 1 Read our strategy and supporting documents to find out what you can do for nature



- 2 Explore our interactive maps to see where you can take the most effective action for nature recovery



- 3 Let us know what you have been doing for nature recovery



[Click here to discover more](#)



Did you know?

Researchers at University College London found that people who spend just two hours a week in nature report significantly better health and wellbeing than those who don't

(White et al., Scientific Reports, 2019).



Strategic aims for nature recovery

The strategic aims for nature recovery in Cambridgeshire and Peterborough are listed below. These have been identified through:



Workshops with farmers, landowners, technical specialists, local authorities, government agencies and environmental organisations

Engagement sessions with communities, interest groups and local people

public surveys

desk based research

Existing strategic environmental reports produced for Cambridgeshire and Peterborough



Air quality


Strategic aim:

Improve the air quality of Cambridgeshire and Peterborough and reduce impacts on habitat and species.

Climate change

Strategic aim:

Strengthen the resilience of Cambridgeshire and Peterborough's environment against the effects of extreme weather and climate change impacts..



Communities

Strategic aim:

Enhance a sense of place where both people and nature can benefit by:

- empowering organisations, communities and interest groups (e.g. faith groups) to manage priority habitats, local freshwater habitats and community greenspaces.
- promoting people's connections to nature and improving health and wellbeing.
- supporting local organisations and interest groups to help recover nature on their doorstep.
- working with residents to identify and improve or provide areas of natural green space close to existing settlements.

Doubling nature

Strategic aim:

Support Natural Cambridgeshire's doubling nature ambition to at least double the area of rich wildlife habitats from approximately 8% to more than 16% by 2040 (and increasing beyond that) and use the Lawton principles to guide nature recovery action:



- **Better** - protection and better management of high-value nature sites
- **Bigger** - increase the size of existing habitats and nature sites
- **More** - creating habitats in line with LNRS priorities
- **Joined Up** - connecting existing and habitats across the landscape



Development

Strategic aim:

Encourage existing, new and large developments to avoid impacts on high-value nature sites, contribute positively to local nature networks, and incorporate wildlife friendly features in gardens, buildings and land by:

- avoiding direct and indirect losses or damage to high-value nature sites
- maximising opportunities to enhance nature through the planning system.
- enhancing and expanding existing greenspaces for nature and providing new strategic green space connected to the wider green infrastructure network by green corridors. This will help to address the lack of accessible green infrastructure, reduce recreational pressure on existing nature sites, provide multifunctional benefits and provide an important asset to meet growing demand from new development.
- introducing urban greening interventions such as green roofs, sustainable drainage systems (SuDS), street trees, wildflower verges, allotments and pocket parks within existing and newly proposed urban areas to deliver multiple benefits for people, wildlife and the environment.
- supporting biodiversity net gain (BNG) policy, encouraging 20% as a minimum, where possible.

Environmentally friendly farming

Strategic aim:

Support farming and food production across Cambridgeshire and Peterborough to be conducted using environmentally sustainable practices, recognising the importance of food production and the potential of farming to support nature recovery.



Geodiversity

Conserve and celebrate Cambridgeshire and Peterborough's rich geodiversity by:

- Conserving the geodiversity character and natural heritage of the rural and urban landscape and the connection to habitats and species.
- Celebrating the links between geodiversity, landscape, and the history and cultural heritage of our villages, towns and cities.

Did you know?

In parts of the Cambridgeshire Fens, peat has started forming at least 6 times (with marine flooding in between) since the last Ice Age - it really does want to grow peat!



Internationally rare chalk streams

Strategic aim:

Recognise the importance of the internationally rare chalk springs and streams within Cambridgeshire and Peterborough and collaborate with partners across sectors to conserve and restore them.



Landscape and Historic environment

Strategic aim:

Celebrate Cambridgeshire and Peterborough's rich culture, built / natural heritage, and its surrounding environment by:


- Understanding the character and distinctiveness of the local landscape to enable the design of nature recovery projects that respond to this and deliver co-benefits that enhance landscape, local distinctiveness and sense of place.
- Protecting the character and integrity of the rural and urban historic environment whilst maximising nature recovery benefits.
- Understanding and promoting protection of the historic environment (e.g. ridge and furrow field systems or historic parks) alongside nature recovery.



Peatland

Strategic aim:

Promote healthy functioning peatlands to store and capture carbon whilst not increasing methane emissions.



Water quality and resources

Strategic aim:

Promote integrated water management techniques considering water resources, water quality, and resilience to flooding to support:

- improvements in water quality by reduction of and better management of pollutants from farmland, transport networks, wastewater works and other rural and urban sources.
- increasing awareness of water scarcity amongst all users
- improving water efficiency for domestic and commercial users implementing options to address water scarcity.
- using nature recovery as part of a natural flood management programme to help mitigate against flood risk.





6

Priority natural landscapes

Priority natural landscapes

Landscape-scale nature recovery

A focus on nature recovery at the landscape scale, looking beyond and connecting individual sites and improving the resilience of the landscape as a whole, has been long recognised as essential in Cambridgeshire and Peterborough. This is particularly the case for an area where around 80% of the land is farmed, most of which is arable crop land, and where high value nature sites are small, fragmented and only cover about 8% of the area. The case for landscape-scale restoration of nature becomes even more important when you consider that Cambridgeshire and Peterborough is one of the fastest growing areas in the country in terms of population growth and new development, and that there aren't large areas of accessible downland, forest, moorland or coast. The impacts from climate change and the urgent need to provide space and connectivity for species to adapt and move further supports this landscape-scale approach.

However, a landscape-scale approach to nature restoration must also align with our

continuing need to grow food and provide the homes, employment sites and other infrastructure we need.



In an intensive agricultural landscape with rapidly growing towns and cities and where high value nature sites are highly fragmented, a landscape-scale approach to nature restoration must be highly focused to be effective in the short-term and to target scarce resources to those areas where the greatest biodiversity benefits can be achieved.

Application of the Lawton principles in Cambridgeshire and Peterborough will look very different to other parts of the country with greater land cover or concentrations of nature sites. Here, the first step is to rebuild our core biodiversity hotspots. We don't have areas with extensive and well-connected habitats such as the Brecks, the Chilterns, or Broads. Most of our best nature sites are either small and / or isolated. Our priorities for nature recovery will therefore look very different to many other parts of the country.

Nature recovery requires us to move beyond a single habitat or species focus. Even within Cambridgeshire and Peterborough there are areas where a greater concentration of nature sites has survived. In many of these there are combinations of remnant habitats whether woodlands, meadows and ponds; or chalk grasslands and scrub; or fens, wet grasslands and wet woodland. These combinations of habitats all in proximity to each other support higher levels of biodiversity than blocks of single habitats.



Did you know?

The Ouse washlands were created 360 years' ago to retain winter flood water from the Ouse.

Identification of priority natural landscapes

In 2000, the Cambridgeshire and Peterborough Biodiversity Partnership identified those areas of Cambridgeshire and Peterborough where the highest concentrations of habitats survived, and which represented the variety of landscapes and habitats across the area.

These were refined by Natural Cambridgeshire into six priority landscapes¹ for landscape-scale nature recovery to deliver the Cambridgeshire and Peterborough doubling nature vision:

Natural Cambridgeshire Priority Landscapes

Each of these areas is briefly described below

Connected Fens

The area of internationally important remnants of ancient fen and washland habitats that now cover less than 1% of the Fens. They include the richest areas of habitat locally, supporting many rare and iconic species. Major restoration is planned and taking place in the Great Fen, Wicken Fen and Ouse Washes areas.




John Clare Countryside:


An area of ancient woodland, species-rich limestone grasslands and wetlands, supporting the richest array of species beyond the fen sites and connecting to the large Rockingham Forest area in Northamptonshire to the west.



Nene Valley

An area of remnant flood meadows and other wetland habitats from the city of Peterborough upstream to the large wetlands along the Northamptonshire Nene.





Great Ouse Valley

An extensive area of floodplain meadows, former sand and gravel pits and other wetlands supporting a rich array of plants and nationally important numbers of waterbirds. It connects the Ouse Washes to the Great Ouse in Bedfordshire.



West Cambridgeshire Hundreds

The ancient woodlands and parklands west of Cambridge to the Bedfordshire border with populations of iconic plants and invertebrates with restricted distributions in England.

Cambridge Nature Network

The area of downland, fen and wooded farmland around Cambridge and the river Cam and its chalk stream tributaries extending into Essex.

While these six landscapes form the top priorities for nature recovery when viewed at the county or regional scale, they are still relatively isolated from each other and similar areas within neighbouring counties. To help create a national nature recovery network these will need to connect with other biodiversity hotspots. Nationally, nature recovery will connect from the English Channel to the north of Scotland and from the Wash to West Wales.


Therefore, landscape-scale nature recovery in Cambridgeshire and Peterborough will also need to take place within other areas between the

six priority landscapes. The Wildlife Trust has worked with district councils and other partners to identify additional priority landscapes to focus nature recovery at the district-scale and as a result, further refined the boundaries of the Natural Cambridgeshire Priority Landscapes.

The detail is set out in reports for the Cambridge Nature Network², East Cambridgeshire Interim Nature Network³, Fenland Interim Nature Network⁴ and Huntingdonshire Interim Nature Network⁵. Separate unpublished mapping studies have been undertaken for the Great Ouse Valley and John Clare Countryside following the same principles.

The aim of these nature network studies was to:

- 1** Identify priority areas for landscape-scale action to support nature's recovery, from both desktop analysis and targeted fieldwork
- 2** Agree the boundaries of the priority landscape areas through stakeholder engagement with key stakeholders and a sample of major landowners
- 3** Identify the critical components of a nature recovery network in each of the priority areas, based on the Lawton Principles



These nature network reports include a detailed analysis of the nature recovery opportunities in each of the priority natural landscapes. The aim within each area is to create or restore at least 30% of the landscape to high value nature sites, with these set within a wider landscape of nature friendly farming. In all but one or two of the priority natural landscapes, farming would continue to be the dominant land use. Creating or restoring land to achieve 30-40% high value nature sites in each of these priority


natural landscape areas would achieve the Cambridgeshire and Peterborough doubling nature vision target for 16% of the Cambridgeshire and Peterborough land area to be high value nature-rich habitats.

Combining the Natural Cambridgeshire six priority landscapes with the district priority natural landscapes produces a coherent, long-term nature network map for Cambridgeshire and Peterborough to support nature recovery at scale.



A long-term nature recovery network for Cambridgeshire and Peterborough

Nature Networks



These priority natural landscape areas provide a focus for delivery of long-term nature recovery in Cambridgeshire and Peterborough and for targeting investment in the natural environment, whether agri-environment schemes, private natural capital finance or other public funds. They also highlight those areas where BNG and offsetting would be best focused.

Within these priority natural landscape areas, partners are undertaking more detailed nature recovery opportunity mapping to identify the best areas for habitat creation and restoration. In some locations, specific areas and sites have been identified and agreed in dialogue with landowners. These have been integrated into the LNRS local habitat map.



Beyond the priority natural landscapes

Although creation of high-value habitats might be focused within priority natural landscapes, there are opportunities for nature recovery across our farmed and urban landscapes. Action to support nature recovery can occur anywhere and be undertaken by anyone.

Creating more nature friendly farmed landscapes and urban areas is an essential part of nature recovery, as it provides space for many species to move between high-value habitats and in response to climate change, and places for people to experience nature close to where they live. As most of our area will remain farmed or be part of the growing urban areas, it is important that these areas are as hospitable to wildlife as possible.

Nature recovery on our farmland

Nature recovery on our farmland

80% of Cambridgeshire and Peterborough is farmed and over 70% of the land is arable. Agriculture will remain the dominant land use across our area. Even in our priority natural landscapes, where increased cover of high value habitats is essential for nature recovery, farming will remain the dominant land use. However, all farms, can play an active role in nature recovery. Many farmers are already doing this, though much more could be done, with the right incentives, encouragement and information.

Did you know?

80% of Cambridgeshire and Peterborough is farmed and over 70% of the land is arable.

A more nature-friendly farmed landscape is essential to support more, bigger and better high-value natural habitats. Nature friendly farming would help provide connectivity between core local nature sites as well as support those species that depend on the farmed environment.


In Cambridgeshire and Peterborough, the widespread adoption of nature friendly farming practices, for example putting unproductive or less productive parts of fields down to field margin habitat options, is an integral part of nature recovery. If the majority of farms were able to put

7-10% of their farm into wildlife friendly options, the decline in many species whether invertebrates, plants or birds could be reversed. While this will not be possible on all farms, even 3-5% cover of field margin options will start to make a difference.

Regenerative forms of agricultural aimed at improving soil health complements nature-friendly farming and will often be undertaken on the same farm. Some farmers will choose an organic approach or adopt agro-forestry principles.

Nature recovery across the farmed landscape will involve a variety of approaches. Most farmers will continue with intensive production but help nature recovery with on-farm actions such as nature-rich field margins, headlands, hedgerows or ditches. Other farmers may transition towards agro-ecological production that prioritises nature recovery and soil health. Some may integrate wildlife-rich grassland habitats with conservation grazing. In a few areas, land may transition to naturalistic grazing or more extensive areas of new grassland, wetland or woodland habitats. All these approaches complement each other at the landscape scale to ensure continued farming and food production plays its part in nature recovery.

Nature friendly farming will take different forms in the different landscapes that make up our area.



The Fens

Farms in The Fens are generally on peat or organic-rich soils with

networks of drainage ditches to manage water. The ditch systems and pumping are used to lower water levels in the fields to allow arable crops to be grown, and at other times to provide water for irrigation during the crop growing season. Much of the land is at or below sea level requiring the water to be pumped into the main drains that take the water to the Wash.

The landscape has few other nature rich habitats beyond the internationally important wetlands, though there are small woods, pockets of grassland, and networks of fen droves. The ditch system provides a connected habitat network across the whole fens landscape. Some parts of this network are rich in aquatic wildlife and relic fen species; however, this is limited to those ditch networks where the water is clean, often arising from underlying gravels, chalk or peat. The water quality in much of the ditch network is poor and supports a low diversity of aquatic species; however, Water Voles are present in nationally important numbers across the internal drainage board (IDB) ditch network, irrespective of water quality.

Climate change and sea level rise with flooding risks from rivers and the sea, combined with the potential for increasing droughts present major challenges to the future of farming. The continued loss of peat with the resultant carbon emissions are also a major challenge. Nature recovery can be part of the response to these challenges.

Drains are conduits and channels providing stepping stones to nature sites between larger nature areas such as the Ouse and Nene Washes, as highlighted in the Great Fen Project. The provision of buffer areas adjacent to the main drains and ditches will assist with the following nature friendly farming options:

- provision of grass or wildflower field margins along ditch networks to reduce agricultural run-off in the ditches and improve water quality. All ditches should have at least a 2-metre margin from the top of the bank.

- management of the ditch network, to create areas that hold water all year round and have a mixture of shallow margins to support wetland plants and steeper sides for Water Vole to burrow into.
- creation of small areas of wet grassland or willows in unproductive field corners.
- creation of small wetlands in strategic locations close to pumping sites or farm reservoirs to filter and clean water and provide small habitat stepping stones across the landscape.
- clusters of farms working together to build farm reservoirs to provide water for farming operations and reduce impacts on water supply elsewhere. These can be designed to include at least one shallow margin suitable for wetland plants and provide habitat for wetland birds and invertebrates.
- adoption of alternative forms of cropping such as wet farming (paludiculture).



The Claylands

The Claylands in the west and south of the area were largely grassland or mixed farming until the 1930s.

However, today arable farming is dominant with small areas of pasture, often on the edge of villages, still present. Areas of species-rich grassland are limited to isolated fields and road verges.

These areas are the most wooded parts of our area and in some parts, there are networks of hedgerows, though many are often thin, gappy and with sparse cover at their base, reducing their value to breeding birds.

Most ponds in the area are within the Claylands and support species such as Great Crested Newt.

The installation of under-soil drainage networks in the twentieth century helped the transition from pasture to arable farming.

However, this, together with straightening and deepening of rivers and streams has contributed to flooding downstream.

Nature friendly farming options for the Claylands include:

- better management of hedgerows to create tall, thick hedgerows, with dense cover at the base and buffers of wide grass or wildflower field margins. Some hedgerows may contain trees.
- restoration and creation of networks of hedgerows to connect clusters of woodland.
- creation of networks of wildflower rich grassy field margins and headlands on less productive or unproductive parts of fields. Incorporation of areas of wild bird seed mixes, pollinator mixes and legume-rich crops.
- creation of wide habitat buffers to watercourses and where possible reconnecting rivers to their floodplains and creating more natural river profiles.
- restoration or creation of ponds, ideally as clusters of two or three ponds.
- natural flood management to slow the flow and improve water quality before it reaches the main rivers. Actions include leaky dams across ditches, creation of small wetlands and water retention basins, planting of hedgerows or grassland buffer strips across slopes, sowing winter cover crops and reducing soil compaction.
- creation of small copses as stepping stone habitats, including tree and woodland planting along ditches and streams as part of natural flood management.
- buffering ancient woodlands with wide flower rich field margins or headlands and allowing some of these to become a mix of flower rich grassland and scrub.



The Chalklands

There is a long history of arable farming across the Chalklands in the south of the area, though mixed farming was the norm before World War Two. Today arable farming is predominant, with only a few mixed arable and livestock farms.

Species-rich grasslands are now restricted to non-farmed land such as ancient monuments, former chalk quarries and road verges.

Rare arable plants occur across the landscape but often associated with farms that include some less intensively cropped areas. Farmland birds associated with open landscapes occur in good numbers where farms include all year-round feeding opportunities.

Fields are generally large and where hedgerows are present, they are usually relatively recent, low growing, and thin with sparse cover at the base.

Small planted farm woodlands occur across parts of the landscape but are rarely more than 100-150 years old. Most are unmanaged and do not support the areas of open space, dense shrubby growth or deadwood that most specialist woodland species depend on.

Nature friendly farming options for the Chalklands include:

- management and creation of varied wildflower rich and grassy field margins across the farmed landscape and adjacent to remnant chalk grassland sites such as road verges or chalk pits.
- creation of cropped but unsprayed headlands and wide field margins for rare arable plants.
- planting of areas with wild bird seed mixes, pollinator mixes and legume-rich crops.
- creation of areas of dense scrub on less productive land across the landscape. Pockets of scrub are likely to better support species such as Turtle Dove than hedgerows in this landscape.
- creation of wide habitat buffers along chalk streams and around spring sources and where possible reconnecting chalk streams to their floodplains.
- creation of more permanent ponds where pockets of clay are present. Water sources are important for wildlife including priority species such as Turtle Dove.
- creation of temporary ponds for specialist species of this rare habitat.
- management of farm woodlands to create more varied woodland habitats including open space, dense shrubby growth and increases in deadwood.

Nature recovery in our cities, towns and villages

Within our cities, towns and villages are networks of public open spaces. Some of these may lie within a priority natural landscape, while others lie outside, but all provide space for people to interact with nature. In some towns and villages, they provide the only access to nature within walking distance for those who live there.

Local councils recognise the environmental importance of their open spaces. Some may be natural greenspaces while others are primarily used for recreation but may have wilder corners. Urban natural greenspaces are often managed with community groups. Parish councils also manage small open spaces under their control. There are significant opportunities to work with local people and community groups to achieve more for nature across our public parks and open spaces e.g. through the development of neighbourhood plans.

Did you know?

You don't have to do everything at once – even small changes can bring big benefits.

There are many private gardens across our towns and villages which potentially provide a haven for urban wildlife from foxes and hedgehogs to garden birds, frogs and insect pollinators such as bees. Everyone can garden for wildlife, whether it is in a detached house with large garden, in a typical modern housing estate small garden or in window boxes in flats. Imagine if 30% of private gardens in Cambridgeshire and Peterborough were managed with wildlife in mind, it would equate to an area slightly larger than the Nene and Ouse Washes combined or similar area to the whole of the Great Fen. The Climate Change Gardening Group has a range of informative tips to help local residents

mitigate against climate change and declines in biodiversity. You don't have to do everything at once – even small changes can bring big benefits.

The public open spaces and gardens provide the basis for the “urban forest” and are supplemented by street trees and road verges. As temperatures continue to rise there is a need for much greater “urban greening” with increased tree and vegetation cover to help provide urban cooling in towns, as well as the other benefits of cleaner air and improved mental health.

Buildings made from locally sourced stones provide habitats for a variety of plants and animals and provide links to our local geodiversity, surrounding landscapes and the historic character of our villages, towns and cities.

Buildings can be made greener through green roofs and green walls. The incorporation of bat and bird bricks or boxes can provide breeding spaces that otherwise wouldn't exist. The use of SuDS is increasing, and these can provide space for wetland wildlife in our towns.

Actions to support wildlife include the introduction of wildflowers, pond restoration and creation, tree and hedge planting, or more wildlife-friendly mowing and hedge cutting regimes. Every parish or community can prepare their own local nature recovery plan, to guide actions in their area, within gardens, the public realm or by the farming and landowning community of their area.

There are a variety of resources such as Local Nature Recovery Toolkit and Biodiversity for All Toolkit that provide ideas and options to help people enhance their local patch. Across town and country, communities working together can play their part in nature's recovery.





7

Priorities for nature recovery

Priorities for nature recovery

Our priority natural landscapes provide a focus for nature recovery action. This is complemented by action to support nature recovery across our farmland and within our cities, towns and villages. However, an LNRS also needs to identify specific habitats and species as local priorities for nature recovery action.

This section identifies the habitats and species we have chosen as our priorities.

The method used to identify these local habitat and species priorities is set out further detail in the Prioritisation and Mapping Methodology document.



Habitats identified as local priorities

Broad habitat	Local habitat priority	
 Woodland	<ul style="list-style-type: none"> ✓ lowland mixed deciduous woodland ✓ wet woodland 	<ul style="list-style-type: none"> ✓ wood pasture and parkland ✓ traditional orchards
 Grassland	<ul style="list-style-type: none"> ✓ lowland calcareous grassland ✓ lowland meadow 	
 Wetland	<ul style="list-style-type: none"> ✓ lowland fen, floodplain wet grassland and associated habitats ✓ reedbed ✓ gravel pits, lakes and reservoirs 	
 Habitat mosaics	<ul style="list-style-type: none"> ✓ mosaic of woodland, grassland and wetland 	
 Rivers, chalk streams and drainage ditches	<ul style="list-style-type: none"> ✓ rivers ✓ chalk streams and springs 	<ul style="list-style-type: none"> ✓ fen main drains ✓ fen drainage ditches
 Urban landscape	<ul style="list-style-type: none"> ✓ urban parks, and natural greenspaces 	
 Farmland landscape	<ul style="list-style-type: none"> ✓ arable field margins ✓ ponds 	<ul style="list-style-type: none"> ✓ temporary pools ✓ hedgerows

Habitat actions

We have developed actions for the priority habitats to support nature recovery.

The method we used to identify these actions can be viewed in our supporting documents (Prioritisation and Mapping Methodology and Habitat Opportunity Modelling Methodology).

Mapped actions are specific, proposed actions designed to deliver the priorities set out in the LNRS by creating or enhancing habitats for nature, like planting trees or restoring wetlands, located in priority areas on the strategy's maps.

Unmapped actions, are potential actions for habitat creation or improvement that are not tied to a specific geographic location on the LNRS maps

but could be applied more broadly or in response to future opportunities.

Supporting actions in an LNRS are activities, beyond direct habitat creation, that support nature recovery and enable delivery of LNRS priorities. These actions have been identified and agreed by stakeholders and can include data collection, stakeholder engagement, policy and project development etc.

Unmapped actions are equally important and necessary to achieve the goals of the LNRS. However, please note that **unmapped actions are not eligible for Biodiversity Net Gain (BNG) strategic uplift.**



For all our priority habitats you can explore our interactive map of opportunity areas

Woodland priorities, actions and supporting actions

Although less than 5% of Cambridgeshire and Peterborough is wooded, our remaining ancient woodlands support species that are scarce elsewhere in the country.

Our ancient woodlands are mainly found in areas with heavy clay soils. These support species such as Oxlip that is only found in Cambridgeshire, Suffolk and Essex and the more widespread iconic and enchanting Bluebell. We also have woods with significant populations of Smooth-leaved Elms, including some species that are either unique to this area or where the majority of their known population exists. Other iconic species include Hazel Dormouse, Nightingale and Barbastelle bat. Our woodlands therefore form a priority for local nature recovery action.

Our wood pasture and parklands support collections of veteran and ancient trees. Pollarded Elm form a component of the veteran and ancient trees present in some of our historic parks and those with populations of these trees are a local priority.

Although we have very little wet woodland, ancient willows form a significant component of the landscape. Given this, and because of the potential for large-scale restoration of wetlands in parts of the Fens and through restoration of mineral sites, wet woodlands have also been chosen as a local priority. They will support a wide range of invertebrates but could also help to recover iconic species such as Black Poplar.

The final type of wooded habitat that has been chosen as a local priority is traditional orchards. Parts of Cambridgeshire and Peterborough still have concentrations of old orchards with veteran fruit trees, including around Wisbech and between Cambridge, St Ives and Ely. These include many local varieties of apple, pear or plum not found elsewhere.



Did you know?

Cambridgeshire is one of only a few places where Grass-poly, an arable plant which needs disturbed ground that's flooded in winter, is found

Theme 1 Woodlands (Wo)

Lowland mixed deciduous woodland

Wo1 ► better manage woodlands to improve resilience and structural diversity

►► ENHANCE Actions

Wo1A – enhance the management of unmanaged woodlands with a minimum size of 1 hectare to UK Forestry Standards to achieve a varied structure and greater diversity.

Wo1B – enhance all woodlands within 200 metres of existing ancient woodlands by managing to UK Forestry Standards to achieve a varied structure and greater diversity and act as corridors, or stepping stones.

►► SUPPORTING Actions

Wo1B – deliver productive forestry and agroforestry, utilising funding opportunities when available, to support the economic and environmental benefits of woodland creation and management, including supporting local timber markets.

Wo1C – monitor and manage the impacts of pests and diseases, by promoting the reporting on outbreaks such as via Tree Alert and following the relevant guidance.

Wo2 ► restore planted ancient woodland sites with locally appropriate trees, shrub and ground layer species

►► ENHANCE Actions

Wo2A – restore designated plantations on ancient woodland sites back to appropriate native species.

Wo3 ► expand and connect existing areas of woodland to increase woodland biodiversity and help combat climate change

►► CREATE Actions

Wo3A – improve biodiversity by creating mixed deciduous woodland consisting of appropriate native or climate change tolerant (European only)

species to increase resilience and diversity. Such newly created woodlands are to be within 2.5 kilometres of existing woodland, though in practice should be much closer (ideally within 500 metres).

Newly created woods would ideally form a woodland block (or group of woodlands within 200 metres of each other) which are at least 40 hectares in size. Woodland creation would be expected over approximately 80% of the site area mapped under this measure, with the remaining 20% a mosaic of other complementary habitats such as species-rich grassland, scrub, ponds, and individual trees

Wo3B – deliver wider environmental benefits through tree and woodland planting in appropriate locations such as upper catchments, adjacent to rivers and within and adjacent to centres of population. Planting in upper catchments and adjacent to rivers helps enhance water quality and reduce flood risk. Planting within and adjacent to urban areas can help to enhance air quality, contribute urban cooling, reduce flood risk and promote health & well-being. Planting sites should ideally be a minimum of 0.5 hectares but may be smaller within urban areas.

►► UNMAPPED Actions

Wo3C – create woodland, scrub and other complementary habitat buffers such as species-rich grassland adjacent to priority woodland sites.

Wo4 ► reduce the impacts of deer within existing woodland to allow for natural regeneration and to improve the woodland biodiversity and economic value of timber

►► SUPPORTING Actions

Wo4A – manage the impacts of deer to allow natural regeneration and structural diversity within existing woodland, reducing the impact on biodiversity and economy. Support the establishment of deer management groups.



Wet woodland

Wo5 ► better manage woodlands to improve resilience and structural diversity

►► ENHANCE Actions

Wo5A– enhance the management of existing wet woodlands to create varied woodland structure at the site and landscape scale, in line with the UK Forestry Standard.

Wo6 ► create areas of wet woodland in suitable locations

►► CREATE Actions

Wo6A – create wet woodland within appropriate hydrological areas, consisting of native or climate change resilient (European only) species. Newly created wet woodland should not be created in areas that buffer internationally designated areas such as the Ouse and

Nene Washes, archaeological sites, or where wet grassland, fen and floodplain meadow habitats have been identified as priorities for habitat creation without further consultation.

►► SUPPORTING Actions

Wo6B – where new planting is being considered, ensure that if peat is present, the Forestry Commission / Natural England Peatland Decision Support Framework is followed so that new woodland benefits peat and fen habitat and does not adversely affect its hydrological or ecosystem functions.

Wood pasture and parkland

Wo7 ► improve the management of ancient and veteran trees within wood pasture and parklands

►► ENHANCE Actions

Wo7A– enhance the biodiversity value of

designated or other mapped parkland and wood pasture sites, ensuring protection of and continuity of veteran and ancient trees in accordance with best practice, and the provision of complementary habitats such as species-rich grasslands, ponds and wetlands.

►► UNMAPPED Actions

Wo7B – improve management of all ancient and veteran trees, including urban / street trees; including creation of an appropriate buffer zone around the trees (root protection zone) and planning for replacement with locally sourced 'plant healthy' stock that is resilient to climate change..

Traditional orchards

Wo8 ► improve the management of traditional orchards and seek to expand them where possible

►► ENHANCE Actions

Wo8A– restore and enhance the biodiversity value of traditional orchards. Where possible ensure such enhancements deliver gains for cultural and landscape continuity and for the genetic diversity of fruit trees. Enhance grasslands under the fruit trees to create complementary species-rich habitats.

►► CREATE Actions

Wo8B– create traditional orchards close to existing traditional orchards. These should deliver enhanced biodiversity and heritage value through the planting of heritage variety fruit trees and the creation of species-rich grassland.



Woodland (all Types)

►► SUPPORTING Actions

Wo9A– identify and map ancient and veteran trees within the landscape.

Wo9B - conserve and enhance ancient, heritage and veteran trees, including pollard willows in river valleys, for their biodiversity and heritage value, planning for the provision of replacement stock and future veteran trees. Put in place measures to protect them and ensure succession such as establishment of root protection,

pruning practice, protection from livestock and nursing of seedlings.



Wo9C - improve management of all trees in and out of woodland, including urban / street trees, planning for replacement with locally sourced 'plant healthy' stock that is resilient to climate change.

Wo9D - ensure integration with Forestry England, local community and local authorities' tree and woodland strategies, as well as funding, to support the creation of new areas of trees and woodlands for public access and enjoyment.

Example of species supported by these habitats

Oxlip



Bluebell



Purple Emperor Butterfly



Hazel Dormouse



Wider benefits



Improved water supply



Timber production



Energy generation



Increased genetic diversity



Inspiration / spiritual values



Tranquillity



Cultural heritage values



Recreation and tourism services



Increased pollination



Disease and pest control



Improved water quality



Improved soil quality



Erosion reduction



Water flow and flood



Climate and carbon storage



Improved air quality



Soil formation



Primary production



Nutrient cycling



Water cycling



Increased biodiversity



[Click here to explore our map](#)

Grassland priorities, actions and supporting actions

Species-rich grasslands are rare in Cambridgeshire and Peterborough, though some of our remaining grassland sites are spectacular. The two main types of grassland present in our area have both been selected as local priorities due to their scarcity.

Lowland calcareous grassland occurs either in the south of Cambridgeshire on chalk soils or west of Peterborough on limestone soils. The chalk grassland in the south is often associated with ancient monuments, though a significant area has survived on the Gallops around Newmarket. These grasslands support a rich array of wildflowers including orchid and insect species. Although subtly different in character, the limestone grasslands are also valuable for a similar suite of flora and insects.

Iconic species include Pasqueflower, Lizard Orchid, and a vast array of insects including Chalk-hill Blue Butterfly, Glow-worm and Large Scabious Mining Bees, Common Lizard and some bird species such as Turtle Dove or Stone Curlew, though these also rely on arable land.

The other major type of grassland is lowland meadow, including lowland flood meadows. The most extensive examples occur in the Great Ouse Valley, where distinctive floodplain meadows are present between St Neots and Needingworth. Beyond the river valley, the remaining flower-rich lowland meadows are small and infrequently scattered across the Claylands, but support wildflowers such as knapweed, Green-winged Orchid and Cowslip.

Theme 2 Grasslands (G)

Chalk and limestone grassland

G1 ► enhance calcareous grassland sites, buffer and enlarge them and create habitat stepping stones and corridors to connect habitats for important flora and invertebrate grassland species

►► ENHANCE Actions

G1A - enhance existing chalk and limestone grassland sites to create a diverse set of micro-habitats to support the diversity of scarce and common species associated with this habitat.

►► CREATE Actions

G1B - improve biodiversity by creating species-rich calcareous grassland adjoining to, and up to 500 metres from, existing designated and other chalk and limestone grasslands. There is no minimum site size, but larger sites are preferable (in combination with other complementary habitats), and there should be a realistic ambition for sites to become



priority habitat in the future (it usually takes 50-60 years to create this priority habitat from former arable land).

Lowland meadows

G2 ► enhance existing meadow and flood meadow sites and create buffers and new sites to provide a refuge for species and improve their resilience to climate change

►► ENHANCE Actions

G2A - enhance existing species-rich neutral grassland sites to support the diversity of scarce and common species associated with this habitat.

►► CREATE Actions

G2B - improve biodiversity by creating species-rich neutral grassland adjoining to, and up to 500 metres from, existing designated and other neutral grassland sites. There is no minimum size threshold, but larger sites are preferable (in combination with other complementary habitats) and there should be a realistic ambition for site to become priority habitat in the future.



Dry grasslands (all Types)

►► CREATE Actions

G3A – deliver wider environmental benefits such as runoff reduction, water quality enhancement and potential access to nature, by creating species-rich grasslands in appropriate locations.

►► SUPPORTING Actions

G4A - encourage the creation of complementary habitats (e.g. scrub, ponds), so long as the

grassland remains the majority habitat created.

G4B – restore limestone quarries to create limestone grasslands, associated habitats and limestone geological features.

G4C - work with ecologists and the local farming sector to develop sustainable grazing solutions for existing and newly created species-rich grasslands.

G4D - establish a local seedbank to be accessible across the area, supplying locally sourced seed mixes seed for different soil types and uses.

Example of species supported by these habitats

Knapweed Broomrape



Waxcap



Glow-worm



Wider benefits



Improved water supply



Timber production



Energy generation



Increased genetic diversity



Inspiration / spiritual values



Tranquillity



Cultural heritage values



Recreation and tourism services



Increased pollination



Disease and pest control



Improved water quality



Improved soil quality



Erosion reduction



Water flow and flood



Climate and carbon storage



Improved air quality



Soil formation



Primary production



Nutrient cycling



Water cycling



Increased biodiversity



[Click here to explore our map](#)



Wetlands priorities, actions and supporting actions

The Fens basin was once the largest wetland in Western Europe, but today less than 0.5% of the area supports wetland habitats. What remains includes the internationally important lowland fen sites of Wicken Fen, Woodwalton Fen, Chippenham Fen and Holme Fen. These are our most species-rich habitats, supporting many rare and scarce plants, invertebrates and other species. These include Crane, Hen Harrier, Spined Loach, Tansy Beetle, Marsh Carpet Moth, Fen Violet, Fen Wood-rush and Greater Water-parsnip.

Our other type of internationally important wetland is the Ouse and Nene Washes. These were created as part of the water management and drainage system that turned The Fens from wetland to farmland. The extensive area of floodplain wet grassland supports internationally important numbers of waterbirds including Bewick's

and Whooper Swan, geese, ducks, wading birds and herons and egrets.

The Fen wetlands are therefore the top priority for nature recovery locally. They could in future provide space for species currently missing from the landscape to be returned such as Spotted Crake, Pool Frog or Swallowtail Butterfly.

Other types of wetland habitat selected as a local priority for recovery include reedbeds and gravel pits, lakes and reservoirs. Reedbeds are scarce across the area, but because of mineral restoration plans, significant new reedbeds will be created including one of the largest in England at Ouse Fen. These support Bittern and Reed Leopard Moth. Former gravel pits and existing reservoirs such as Grafham Water have been chosen as a local priority because they support nationally important numbers of wetland birds.

Theme 3 Wetlands (We)

Lowland fens and floodplain wet grasslands and associated habitats

We1 ► create wetlands on peat around remnant fen sites to buffer and protect the peatland fen habitats for their special biodiversity and improve their hydrological function

►► CREATE Actions

We1A – in the following large-scale Fenland wetland vision areas: Wicken Fen; Great Fen; Ouse Washes landscape area; and adjacent to the Nene

Washes, create mosaics of wetland habitats such as fens, reedbeds, wet grasslands for breeding and wintering waterbirds, wet woodlands, pools and ponds on peat and other suitable soils. On the higher, drier ground within these vision areas on non-peat soils, create other complementary habitats such as meadows, scrub, woodland and ponds.

We2 ► identify and prioritise areas for the protection of peat to restore fen habitats and reduce carbon losses from the soil

►► ENHANCE Actions

We3A – manage floodplain wetland mosaics through sustainable grazing to create habitats that are wetter for longer and support a greater diversity and abundance of wetland flora and fauna, including breeding and wintering wetland birds.

We3 ► enhance and create floodplain wet grassland sites to better support breeding and wintering wetland birds

►► CREATE Actions

We3B – create areas of floodplain wetland mosaics within the embanked floodplains of the major rivers and other suitable locations to enhance connectivity between wetland sites and reverse habitat fragmentation to support diverse species assemblages

Reedbeds

We4 ► sustainably manage established reedbeds to preserve their important role as a valuable habitat and carbon sink

►► ENHANCE Actions

We4A – enhance existing reedbeds to benefit the bird and invertebrate species associated with this habitat. This can include managing reeds to a variety of heights at different stages of growth, areas of wet reedbed with varying water depths, dry reedbed and a network of open water pools and channels.

►► CREATE Actions

We4B – create reedbed habitats that are resilient to climate change in suitable locations beyond the landscape-scale wetland vision areas. Large examples are likely to be associated with the

restoration of mineral sites.

Gravel pits, lakes and reservoirs

We5 ► better manage gravel pits to support wetland wildlife

►► ENHANCE Actions

We5A – enhance existing former sand and gravel pits sites through best practice management to maintain areas of open water, fringing reedbeds and other wetland habitats such as pollard willows and wet woodland for wetland birds, invertebrates and flora. Reduce nutrient enrichment and pollution and where necessary reprofile edges to create shallow sloping margins and bare areas for natural regeneration.

We6 ► design new mineral sites and reservoirs to include features suitable for wetland wildlife and as stepping stone habitats for wetland species

►► CREATE Actions

We6A – create a range of wetland habitats including wet grassland, reedbeds and ponds around the Fens reservoir.

►► UNMAPPED Actions

We6B – create new bodies of open water after mineral extraction has taken place. Ensure they are designed to promote nature in accordance with best practice, and include shallow margins, islands, areas of bare ground, fringing reedbeds and areas of wet woodland, pollard willows and geodiversity features.

►► SUPPORTING ACTIONS

We6C – Design the Fens reservoir to include, where possible, varied, biodiverse margins and provide space for waterbirds to feed and rest.





We6D– promote the creation of nature-friendly farm reservoirs to capture water for irrigation uses, reduce demand on public water supply, and provide wetland stepping stones across the landscape.

Wetlands (all types)

We7 ► work with farmers, landowners and internal drainage boards to deliver nature-rich habitat restoration and more sustainable land use practices to protect peat, through the adoption of new agri-environment schemes and minimising the presence of bare peat

►► SUPPORTING Actions

We7A – enhance water management systems and manage land connected to peatland habitats to protect their hydrological and nutrient status, for example reducing nutrient rich run off and modifying drainage next to lowland fens to reduce or prevent water losses from the site.

We7B – work in partnership with key stakeholders to pursue sustainable long-term water management solutions, ensuring water is captured,

stored and conveyed to the correct places at the correct time, through an Integrated Water Management approach that looks at site, farm and landscape scale to support creation and enhance of wetland habitats.



We7C– farm businesses should help reduce the effects of agricultural run-off on water quality including implementation of integrated pest management and reduction of pesticide use, and best practice nutrient management to reduce nutrients entering watercourses.

We7D – Promote farming and land use changes within areas of lowland peat, as identified on the England Peat Map <https://england-peat-map-portal-ncea.hub.arcgis.com/>,


that significantly reduce peat losses and carbon emissions and contribute to reduced pollution and improved water quality. In most cases this is likely to involve alternative farming practices or crops, which may include novel and emerging approaches such as paludiculture or other forms of wet farming. In some locations this may involve wetland and other habitat creation.

We7E– promote the creation of nature-friendly farm reservoirs within the Fens to capture water for irrigation uses, reduce demand on public water supply, whilst providing wetland stepping stones across the landscape.

We7F – continue to map areas of surface and buried peat to better inform peat conservation and restoration and support the use of peatland maps by

partners and stakeholders.

We7G – increase educational opportunities across the Fens landscape to access its geodiversity, archaeology and cultural heritage to enhance enjoyment and understanding for those who live and work in and visit the Fens, supporting the development of tourism opportunities to benefit the area's economy and communities.



Example of species supported by these habitats

Ground Beetle



Fen Violet



Yellow Iris



Drinker moth caterpillar



Marsh Fragrant Orchid



Wider benefits



Improved water supply



Timber production



Energy generation



Increased genetic diversity



Inspiration / spiritual values



Tranquillity



Cultural heritage values



Recreation and tourism services



Increased pollination



Disease and pest control



Improved water quality



Improved soil quality



Erosion reduction



Water flow and flood



Climate and carbon storage



Improved air quality



Soil formation



Primary production



Nutrient cycling



Water cycling



Increased biodiversity



[Click here to explore our map](#)


Rivers, streams and drains priorities, actions and supporting actions

All rivers and their floodplains have been selected as local priorities for nature recovery as they provide landscape-scale corridors and can deliver a wide range of other natural benefits. The upper Cam catchment supports around a dozen chalk streams, although many have been significantly modified by river engineering, drainage and excessive water abstraction for human use. Iconic species include Brown Trout and Water Vole. England has most of the world's chalk streams so even damaged examples are a priority for nature recovery.

Our other rivers support species such as Otter, Kingfisher and populations of fish and aquatic insects such as dragonflies. Some species such as Brook Lamprey are only associated with unpolluted water. In the medium to long term there may be opportunities to re-introduce beavers to some tributaries and upper catchments of the Cam, Great Ouse and Nene, to help deliver nature-based solutions to reduce flood risk and improve water quality. However, throughout most of our

area this could not be considered before significant riverside habitat, and woodland creation has been undertaken to create a suitable environment.

The network of fen main drains and drainage ditches is essential to continued farming of the Fens, particularly with much of the land at or below sea level. Although man-made, the drainage network supports relic fen plant and invertebrate species, particularly where the water quality is good. Even where the water quality is poorer, the network supports a nationally significant population of Water Vole and is also home to the rapidly declining European Eel. They therefore provide an interconnected wetland network across the landscape, that can support the movement of species between the major wetland restoration sites.



Theme 4 Rivers, streams and drains (RD)

Rivers

RD1 ► create better and wider buffers alongside rivers and streams to enhance the water quality and biodiversity of our rivers

►► ENHANCE Actions

RD1A– establish natural or semi-natural buffer zones 50 metres wide adjacent to all rivers and streams to improve river water quality, and consequently their biodiversity value and ecological functionality. Within such buffer zones, appropriate measures should be introduced which both enhance biodiversity and reduce sediment or pollutant run off into the water. Restore the biodiversity and natural function of the river / stream through restoration of the channel, reconnecting rivers with floodplains and creating backwaters and backchannels to provide fish spawning and riparian habitats along the river.

RD2 ► restore natural features within rivers and floodplains to provide a wide range of benefits for the environment including reduced flood risk and improved water quality

►► UNMAPPED Actions

RD2A– reduce sedimentation of rivers by better land management of the river catchment to create habitats that help prevent soil erosion and stop sediments entering watercourses.

Chalk streams

RD3 ► conserve and restore chalk streams, their water quality and associated habitats to preserve and improve their high ecological value to both nature and people

►► ENHANCE Actions

RD3A– establish natural or semi-natural buffer zones up to 50 metres wide adjacent to all chalk streams and their headwaters, winterbourne sections and springs to improve river water quality, and consequently their biodiversity value and ecological functionality.

This will help reduce sedimentation of rivers by better land management of the chalk river catchment to create habitats that help prevent soil erosion and stop sediments entering watercourses.

It will also help prevent spray drift from agricultural activities into the chalk streams.

Rivers and streams (All Types)

►► SUPPORTING Actions

RD6A – reconnect rivers to floodplains wherever feasible to hold more water for longer on floodplains to provide more space for wetland biodiversity, and to contribute to nature based solutions such as improved water quality, reduced flood risk, and water resources.

RD6B – identify and remove redundant artificial structures within rivers and streams to create more natural flows and remove barriers to fish migration. Where barriers cannot be removed (e.g. navigation locks), seek to maximise the habitat variety within backwaters and bypass channels.

RD6C – continue to advocate for early reductions in abstraction from the chalk aquifer to restore natural flows to chalk springs and streams.

RD6D – undertake definitive mapping of chalk streams and their headwaters and springs to inform water resource planning, water quality improvements and future chalk stream restoration efforts, working with partner organisations (e.g. Natural England, Environment Agency, Cam Catchment Partnership and Greater Cambridge Chalk Stream Project).

RD6E – continue to support existing farm clusters, working with catchment partnerships and local facilitators to deliver high quality chalk streams, and address issues around water quality and water scarcity by supporting alternative water storage solutions and aquifer recharge.

RD6F – work in partnership with water companies, catchment partnerships, landowners and citizen science projects to deliver comprehensive monitoring of the chalk streams, including headwaters and springs.

RD6G – work in conjunction with organisations already undertaking invasive species control to eradicate mink and other invasive species. This should be carried out with careful consideration of ecological impacts and in compliance with relevant regulations and guidelines.

RD6H – enhance sections of watercourses

identified as bathing spots for the benefit of both biodiversity and health and wellbeing.

RD6I – Promote understanding of the flow of rivers across our landscapes, connecting our villages, towns and cities to the countryside and, eventually, to the sea, along with their natural evolution over time and the accompanying record of environmental change.



Fen main drains

RD4 ► enhance the fen main drains and their adjacent land to create wetland stepping stones along these landscape corridors

►► ENHANCE Actions

RD4A – maximise opportunities for nature in fen main drains by choosing nature-friendly maintenance and management approaches such as enhancing in channel habitats to provide more habitat for wetland wildlife alongside managing flood risk and water levels.

►► CREATE Actions

RD4B – within corridors 50 metres either side of the fen main drains, create wetland stepping-stone habitats including reedbeds, fens, wet grassland, ponds and open water features (taking into account the 20 metre IDB bylaw maintenance corridors).

►► UNMAPPED Actions

RD4C – support aquatic nature in fen main drains by reducing or intercepting direct discharges of nutrient-rich water into fen internal drainage board ditches.

Fen drainage ditches

RD5 ► enhance the network of fenland farm drainage ditches by providing habitat buffers and improving water quality

►► ENHANCE Actions

RD5A – maximise opportunities for nature in fen drainage ditches by choosing nature-supporting and enhancing maintenance and management approaches such as enhancing in channel habitats

to provide more habitat for wetland wildlife alongside managing flood risk and water levels.

►► CREATE Actions

RD5B– establish uncultivated riparian buffer zones at least 15 metres wide from the top of the bank (including the 9 metre internal drainage board bylaw maintenance corridor). Plant or promote low growing, non-woody vegetation such as diverse grasses or wet grassland mixes that are compatible with regular ditch maintenance operations, to provide habitats for nature

►► UNMAPPED Actions

RD5C– improve water quality monitoring across internal drainage board ditches to identify those



that have poor water quality and develop natural in-channel or off-line filtration areas such as reedbeds, to intercept and remove excess nutrients from water.

RD5D– enhance the network of fen ditches and farm drainage ditches through the provision of buffer zones (a minimum of two metres in width, from the top of the bank). Within such buffer zones, appropriate measures should be introduced which both enhance biodiversity and reduce sediment or pollutant run-off into the ditch.

RD5E–maximise the opportunities for aquatic biodiversity by improving water level management across smaller areas, enabling some seasonally dry ditches to retain water all year round.

Example of species supported by these habitats

Spined Loach



Opposite-leaved Pondweed



River snail



Wider benefits



Improved water supply



Timber production



Energy generation



Increased genetic diversity



Inspiration / spiritual values



Tranquillity



Cultural heritage values



Recreation and tourism services



Increased pollination



Disease and pest control



Improved water quality



Improved soil quality



Erosion reduction



Water flow and flood



Climate and carbon storage



Improved air quality



Soil formation



Primary production



Nutrient cycling



Water cycling



Increased biodiversity



[Click here to explore our map](#)



Habitat mosaics priorities, actions and supporting actions

Many species depend on more than one habitat type to complete their life cycle. Our lowland landscapes also comprise a mixture of wooded, grassy or wet habitats and a combination of these will often result in “the whole being more than the sum of the parts”.

Nature recovery across Cambridgeshire and Peterborough will require the rebuilding of nature

networks comprising mixtures and mosaics of different habitat types and geodiversity features to support the widest range of biodiversity at both the landscape and individual site-scale.

In recognition of our local circumstances, habitat mosaics have been chosen as an additional local priority.

Theme 5 Habitat mosaics (M)

Habitat mosaic

M1 ► improve the diversity of habitats at a landscape and site-scale to increase the diversity and abundance of species and make them more resilient to climate change

►► CREATE Actions

M1A - where two or more mapped actions arise on the same land parcel, the preference is to create mosaics of the identified habitats, with at least one of the habitats of a sufficient quality to meet the minimum standards for that habitat.

►► UNMAPPED Actions

M1B – within the following priority natural landscape areas, create the habitat as described in

the applicable brackets for each area:

John Clare Countryside (woodland, scrub, wood pasture, limestone grassland, lowland meadows, ponds and networks of hedgerows).

West Cambridgeshire Hundreds (woodland, scrub, wood pasture, lowland meadows, ponds and networks of hedgerows).

Cambridge Nature Network – Boulder Clay and Woodlands (woodland, scrub, lowland meadows, ponds and networks of hedgerows).

East Cambridgeshire Nature Network – Boulder Clay Woodlands (woodland, scrub, wood pasture, lowland meadows, ponds and networks of hedgerows).

East Cambridgeshire Nature Network – Soham Grasslands (lowland meadows, scrub, ponds and networks of hedgerows).

East Cambridgeshire Nature Network

– Chettisham to Ely North (lowland meadows, scrub, ponds and networks of hedgerows).

East Cambridgeshire Nature Network

– Breckland Edge (woodland, scrub, wood pasture, lowland meadows, ponds and acid grasslands).

Huntingdonshire Nature Network

– Grafham-Brampton-River Kym (woodland, scrub, wood pasture, lowland meadows, ponds and networks of hedgerows).

Huntingdonshire Nature Network – Holme-Conington (woodland, scrub, wood pasture, lowland meadows, ponds and networks of hedgerows).

Huntingdonshire Nature Network – Folksworth-

Etton-Ashton Wold (woodland, scrub, wood pasture, lowland meadows, ponds and networks of hedgerows).

Huntingdonshire Nature Network

– Hail Weston to Bushmead (woodland, scrub, lowland meadows, ponds and networks of hedgerows).

M1C - restore and enhance habitats within and around SSSIs and County Wildlife Sites, in coordination with landowners, to increase the ecological value of the sites.



SUPPORTING Actions


M1D – work with landowners to identify and map the locations within each priority natural landscape area that can be taken forward to deliver the desired landscape-scale mosaic of habitats and geodiversity features.

Example of species supported by these habitats

Skylark



Common Lizard



Gatekeeper Butterfly



Wider benefits



Improved water supply



Timber production



Energy generation



Increased genetic diversity



Inspiration / spiritual values



Tranquillity



Cultural heritage values



Recreation and tourism services



Increased pollination



Disease and pest control



Improved water quality



Improved soil quality



Erosion reduction



Water flow and flood



Climate and carbon storage



Improved air quality



Soil formation



Primary production



Nutrient cycling



Water cycling



Increased biodiversity




[Click here to explore our map](#)

Urban landscape priorities, actions and supporting actions

Most people in Cambridgeshire and Peterborough live in urban areas whether towns, cities or villages.

Contact with nature improves quality of life and provides many health benefits whether it is the sounds of bird song, the colour of flowers and butterflies, or an urban landscape with good tree cover. Although not one



habitat type, we recognise the value of urban parks and greenspaces and private gardens to human endeavours and quality of life and have therefore included them as a local priority, to bring nature recovery close to people.

Our urban areas support several iconic species including Swift, Swallow, House Martin, Song Thrush, Hedgehog and Common Carder Bee.

Theme 6 Urban landscapes (U)



Urban parks and natural greenspaces

U1 ► enhance greenspaces in urban and rural areas to increase access to nature, foster improved health and wellbeing, and improve the resilience of people and nature to climate change

►► ENHANCE Actions

U1A - enhance the biodiversity value of parks and green spaces within our cities, towns and villages including country parks, public parks and gardens, community gardens and river corridors.

U2 ► create new natural greenspaces in urban areas to provide environmental benefits such as improved air quality, climate change resilience, and greater health and wellbeing for residents

►► SUPPORTING Actions

U2B - increase the urban tree canopy by planting native and climate resilient tree species in streets, parks, and other public spaces to provide habitat, help with urban cooling, and improve air quality.

►► CREATE Actions

U2A - create new natural greenspaces prioritising areas at most risk of health impacts (as identified by the Environmental Justice Index). New natural greenspaces would be of a minimum 0.5 hectares in size, with larger sites preferable, to contribute to natural greenspace accessibility standards. Sites already planned include Whittlesey Country Park, West End Park extension in March, and Ely North Country Park.

►► UNMAPPED Actions

U2C - identify and bring forward through local plans, locations for provision of new strategic natural greenspaces for the market towns in the Fens, including Ely and Littleport in East Cambridgeshire and the locations identified at Chatteris, March, Whittlesey and Wisbech through the Interim Fenland Nature Recovery Network

U2D - enhance churchyards and cemeteries to promote nature friendly management so they provide places for quiet contemplation where wildlife can be seen and heard.

U2E - identify priority Open Mosaic Habitats (nature-rich brownfield



sites) and ensure that their constituent species are conserved and provided for as part of local plan policies, development design and planning decisions.

Transport corridors

U3 ► increase wildlife connectivity across transport corridors

►► CREATE Actions

U3A- connect fragmented habitats through the delivery of strategic wildlife crossings of major road and rail infrastructure (e.g. green bridges to support landscape connectivity and nature recovery). New major infrastructure projects such as East West Rail should include green bridges as a standard part of their design.

U4 ► increase wildlife habitats along infrastructure routes to improve people's access to nature and foster their health and wellbeing

►► UNMAPPED Actions

U4A- support high quality wildlife corridors and pollinator routes along the road and rail network, public rights of way, and ancient routeways such as fen droves.

U4B- map the locations of fen droves and identify remaining biodiversity hotspots, including for locally significant species such as elms and scarce invertebrate species.

Urban greenspaces (all types)

►► SUPPORTING Actions

U5A - use Natural England Green Infrastructure Standards to identify natural greenspace deficits

in all communities across Cambridgeshire and Peterborough. Through local plans, put in place policies and site allocations to address identified deficiencies.

U5B- enable increased access for people of all abilities to appreciate, engage, enjoy and be involved with nature close to where they live and work (e.g. greenspaces, allotments, churchyards, gardens).

U5C - support local communities to develop their own local nature recovery plans that identify, protect and restore sites, habitats and species found in their local area.

U5D - encourage the adoption of nature friendly gardening in private gardens through campaigns to signpost people to toolkits and other online guides.

U5E - identify nature sites adversely impacted by recreational pressures and where necessary plan for and implement suitable alternative natural green space provision, visitor management strategies and implementation of measures such as zoning or closures at impacted sites.

U5F - implement best practice in green infrastructure design and provision in new developments and through local plans.

Promote the retrofitting of urban greening and nature-based SuDS measures where appropriate to do so..

U5G- support communities to explore the built fabric of our villages, towns and cities, connecting people to the local geodiversity and providing new



opportunities to engage with nature alongside our own histories and culture in an ever-changing world.

U5H – enhance the skills base for nature recovery by addressing the shortage of professionals in

habitat restoration, conservation contracting, and species management. By increasing training opportunities for botanists, ecologists, and specialists in species identification, translocation, and reintroductions to ensure effective conservation efforts.

Example of species supported by these habitats

Hedgehog



Brimstone butterfly



Swallow



Bee Orchid



Wider benefits



Improved water supply



Timber production



Energy generation



Increased genetic diversity



Inspiration / spiritual values



Tranquillity



Cultural heritage values



Recreation and tourism services



Increased pollination



Disease and pest control



Improved water quality



Improved soil quality



Erosion reduction



Water flow and flood



Climate and carbon storage



Improved air quality



Soil formation



Primary production



Nutrient cycling



Water cycling



Increased biodiversity



[Click here to explore our map](#)

Farmed landscape priorities, actions and supporting actions

All farms can support nature recovery with the provision of a variety of field edge habitats on less productive parts of the farm. Nature friendly farming is essential to support the habitat creation required to drive nature recovery.

However, there are some species which depend on farming and in-field or boundary habitats. Across our area there are populations of rare and scarce plant species associated with arable crops. There is also a group of birds associated with farmland where we can contribute towards their national recovery, including Turtle Dove, Grey Partridge, Yellow Wagtail, Corn Bunting, Skylark, Yellowhammer and Stone Curlew. East Anglia is also a national stronghold for Brown Hare. These rare arable plant and farmland bird assemblages are a local priority.

Although not renowned for our wooded and hedgerow landscapes there are areas, including on the clay soils and the fen-edge droves, that

support our rare and scarce Elm species and the insect species that are dependent on them. These have therefore also been chosen as a local priority.

While the number of ponds has been much reduced over the past century, they support species such as Great Crested Newt. The UK is the world stronghold for this species, giving us added responsibility for its future. Ponds are essential to support a wide range of wildlife and complement other habitat creation, providing habitats for aquatic plants, insects and amphibians and drinking water for other fauna. Pond restoration and creation can occur across farmland and within our urban areas wherever there are suitable soils, so have been chosen as a local priority.

Temporary pools within chalk farmland support a highly specialised set of plants and invertebrates and some notable examples are known from Cambridgeshire. These have therefore been selected as a local priority.

Theme 7 Farmed landscapes (F)

Arable field margins

F1 ► increase the presence and quality of biodiversity rich arable field margins and headlands around farmland to improve the diversity of farmland species and provide habitat that is connected and more resilient to climate change

►► UNMAPPED Actions

F1A – identify hotspots of rare arable plant species and promote suitable stewardship in field and margin measures in the areas around these hotspots.

F1B – identify hotspots for declining farmland bird assemblages and promote a suite of suitable Stewardship measures in the areas around these hotspots.

►► SUPPORTING Actions

F1C – promote nature-friendly farming where at least 7-10% of every farm is a mixture of high-value habitats, boundary habitats and / or uncropped and unsprayed land including field margins and headlands, pollen and nectar mixes and wild bird mixes.

F1D – encourage farmers to follow best practice to reduce the effects of agricultural runoff including pesticides and excess nutrients on water quality. Measures include implementation of integrated pest management and use of suitable Environmental Land Management options to buffer high-value habitats and watercourses.

F1E – develop a toolkit to promote appropriate Environmental Land Management options for the three main landscape areas of the area – the Fens, the Chalklands and the Clayland.



Farm ponds

F2 ► increase the number of farm ponds, including the restoration of historic pools, to provide important stepping-stone habitats and water supply for wildlife

►► UNMAPPED Actions

F2A – manage networks of farm ponds to support aquatic biodiversity as well as provide a valuable water source for terrestrial wildlife.

F2B – restore historic ponds that have been previously infilled (ghost ponds).

Ponds should be surrounded by uncropped and unsprayed buffers to prevent pollution.

F2C – create networks of new farm ponds across the farmed landscape, particularly on clay soils. Prioritise locations that provide connection / stepping stones to existing areas of priority habitat and wildlife sites. Creation of clusters of two or three ponds is better than single ponds. Ponds should be surrounded by suitable uncropped and unsprayed buffer zones



or other habitat types to prevent pollution. Ponds should be created following best practice design.

Temporary ponds / pools

F3 ► increase the presence of shallow pools on chalk soils

►► UNMAPPED Actions

F3A– identify the best locations to create temporary water bodies and pools in chalk landscape locations known to support species associated with this scarce habitat type. Avoiding ploughing and spraying of seasonally wet areas, and do not create permanent ponds.

Hedgerows

F4 ► increase the presence of hedgerows around farmland to control run-off, improve the abundance and diversity of farmland species and provide habitat that is connected and more resilient to climate change

►► UNMAPPED Actions

F4A– enhance the biodiversity value of existing hedgerows to maximise their wildlife benefits through the implementation of best practice hedgerow management.

F4B – identify all fen droves with important populations of elms or assemblages of invertebrate species and prioritise the protection and enhancement of the hedgerows and associated grasslands.

F4C – plant new native hedgerows to create ecological links between two existing features such as woodlands, priority hedgerows and historical hedgerows.


Where such new hedgerows are created, they should be buffered by strips of



uncultivated land at least six metres wide adjacent to at least one side of the hedge created.

F4D– create networks of hedgerows between core local nature sites within the priority natural landscape areas on clay soils, as listed below:

- **West Cambridgeshire Hundreds**
- **Cambridge Nature Network:** Boulder Clay and Woodlands
- **East Cambridgeshire Nature Network:** Boulder Clay Woodlands; Soham Grasslands; Chettisham to Ely North



- **Huntingdonshire Nature Network:** Grafham-Brampton-River Kym; Fen-edge Woodlands; Folksworth-Etton-Ashton Wold; and Hail Weston to Bushmead

►► SUPPORTING Actions

F4E– encourage the planting of species appropriate to the local area, selected for the soil types, resilient to climate change and supporting local invertebrate species. Specifically for Cambridgeshire, include local varieties of elm to increase disease resilience maintain local genetic variety and provide food source for invertebrates. Avoid the use of non-native species or invasive species.

Example of species supported by these habitats

Harvest Mouse



Poppies



Corn marigold



Broad-leaved spurge



Wider benefits



Improved water supply



Timber production



Energy generation



Increased genetic diversity



Inspiration / spiritual values



Tranquillity



Cultural heritage values



Recreation and tourism services



Increased pollination



Disease and pest control



Improved water quality



Improved soil quality



Erosion reduction



Water flow and flood



Climate and carbon storage



Improved air quality



Soil formation



Primary production



Nutrient cycling



Water cycling



Increased biodiversity



[Click here to explore our map](#)

Site-specific actions

★ Priority natural landscape ► John Clare Countryside

Actions

- Around Castor Hanglands NNR, create a landscape mosaic of habitats with approximately 40% woodland, 20% scrub and 40% lowland meadow.
- Around Barnack Hills and Holes NNR and connecting to Walcot Park, the Southorpe SSSIs / CWS and Burghley Park, seek to create 80% limestone grassland with 20% scrub and networks of hedgerows.
- Along the Whitewater Brook create a mosaic of fens, wet and dry species-rich grassland types, scrub, woodland and wet woodland in roughly equal proportions, and restore a more natural channel.
- Along the Maxey Cut habitat corridor create a mixture of 50% grassland, 20% wetland and 30% woodland associated with gravel extraction sites.
- Adjacent to Bedford Purlieus NNR, create a mosaic of habitats within the former Thornaugh Quarry, including open mosaic habitats, species-rich grassland, scrub, woodland, ponds and limestone rock faces.



★ Priority natural landscape ► West Cambridgeshire Hundreds

Actions

- Create new woodland (60%) and associated habitats (lowland meadow 20%, scrub 20%, ponds and wood pasture) adjacent to selected SSSI woodlands and as stepping stone habitats in locations identified with landowners.

★ Priority natural landscape ► Great Ouse Valley

Actions

- Wildlife Trust detailed field scale mapping to be used ahead of habitat modelling outputs.





Huntingdonshire Nature Network Holme to Conington

Actions

- In the area identified west of Holme Fen, create fen, wet grassland, wet woodland and other complementary habitats to buffer and extend Holme Fen National Nature Reserve.



Whittlesford – Thriplow Hummocky Fields Extension

Actions

Create an arable landscape sensitively managed for the benefit of rare arable flora comprising regularly disturbed soils, a network of ephemeral pools and reinstated historic ponds and avoidance of use of herbicides and fertilisers, to allow opportunities for expansion of flora populations found within the neighbouring SSSIs.



Cambridge Nature Network ► Gog Magog Hills

Actions

- Adjacent to and connecting Cherry Hinton Chalk Pits, Roman Road and Gog Magog Golf Course SSSI create a network of chalk grassland (80%) and associated habitats e.g. scrub and bare ground (20%).
- Create a landscape mosaic of chalk grassland (60%), lowland meadow (20%), fen (10%) and scrub (10%) adjacent to and connecting Fleam Dyke and Fulbourn Fen SSSIs.
- Create a landscape mosaic of chalk grassland (60%), neutral grassland (20%), scrub (10%) and woodland (10%) at Lower Valley Farm habitat bank adjacent to Roman Road SSSI.
- Adjacent to Wandlebury Country Park and Magog Down CWS create a network of chalk grassland (80%) and associated habitats e.g. scrub and woodland (20%).
- Create chalk grassland as part of the new Stapleford Country Park.



Cambridge Nature Network ► Cambridge Fens

Actions


- Adjacent to and connecting Wilbraham Fen, Great Wilbraham Common and Fulbourn Fen SSSIs, create a landscape mosaic of fen (20%), reedbed (10%), wet grassland (40%), lowland meadow (20%), and scrub (10%) habitats.
- Restoration and buffering of Little Wilbraham River and headwaters with a mosaic of species-rich grassland (60%), scrub (20%) and woodland (20%).



Cambridge Nature Network ► Boulder Clay and Woodlands

Actions

- Create new woodland (60%) and associated habitats (lowland meadow 20%, scrub 20%, ponds and wood pasture) adjacent to selected SSSI woodland and at a small number of stepping stone habitat locations identified by landowners.





East Cambridgeshire Nature Network ► Chippenham Fen

Actions

- Adjacent to and connecting Chippenham Fen, Snailwell Meadows and Fordham Woods SSSIs create a mosaic of fen (20%), lowland meadow (50%), scrub (10%) and woodland (20%) habitats.



East Cambridgeshire Nature Network ► Newmarket Chalk Grasslands

Actions

- Create an extended buffer of chalk grassland (80%) adjacent to Devil's Dyke SSSI with up to 20% scrub and bare ground habitats.



[Click here to explore our map](#)



Species identified as local priorities

In addition to habitat priorities, this strategy has identified species priorities. We have a list of 187 species identified as locally and nationally important by both expert stakeholders and the wider public. The methodology for selecting these species can be read in Supporting Document 1 - Prioritisation and Mapping Methodology.

Cambridgeshire and Peterborough holds a large proportion of the British population for some plant species and select animal groups, which therefore hold special importance. Most 'special species' are reliant on habitats which are themselves of special value in the area. About half the species are associated with fenland, either ancient fens or species-rich fen ditches. Ancient woodland and chalk grasslands are also of particular importance. Conservation action for the habitats should meet the needs of most of these species - they can be seen as among the better indicators of the success of habitat conservation.

However, for some local priority species, general habitat actions may not be sufficient and therefore specific actions to support their recovery have been identified.

The priority species identified have therefore been divided into two main groups. Those with actions listed (in this part of the report) and those without

actions listed (see document Supporting Document 5A - Priority Species without actions). Specific targeted conservation actions are still needed and will help many of the species listed without actions (as well as general habitat actions) however detailing these is outside the scope of this LNRS.

For the priority species with actions listed, the majority are species currently present in the area which have been deemed to have a high threat of potential extinction at the local Cambridgeshire and Peterborough level due to being present at only a few sites in the area or for other reasons which make them vulnerable to steep population declines.

Some at the end of this section have been identified as suitable for potential translocation programmes, see - 'Priority species with actions for translocation', and others for potential re-introduction, see - 'Priority species with actions (wild ambition species)'.

For the complete list of priority species please see Supporting Document 5B Full Priority Species List

Many people (including experts and members of the public) helped contribute information for identifying the priority species, including the reasons for inclusion and the potential actions. Thank you to all those who contributed.

Species actions



The following section outlines the actions required to help our priority species recover, where this will not be delivered through the habitat actions.

Black-Tailed Godwit *Limosa limosa*

►► Actions

ACTION 1 ► maintain optimal water levels of wetland habitats (less than 10cm deep) during breeding season.


ACTION 2 ► implement predator control measures, including fencing.

ACTION 3 ► create new wetland habitat (that is not subject to seasonal spring - summer flooding)

ACTION 4 ► support and help deliver the 2023-2033 National Action Plan by working with the existing Project Godwit partnership (<https://projectgodwit.org.uk/wp-content/uploads/2023/09/National-Action-Plan-for-BTG-2023-2033.pdf>)

Justification for inclusion

The UK breeding population is largely concentrated



Black-Tailed Godwit

(90%) at the Ouse and Nene Washes. The overall population suffered declines between 2006 and 2017 (from 48 pairs to 35) but increased to 49 pairs in 2020. The population at the Purls Bridge Pilot Project, Ouse Washes, continue to increase, with seven breeding pairs in 2020 (up from four in 2019 and two in 2018). But there is still no breeding on the Ouse Washes themselves and, at both reserves, the populations are cushioned by captive reared head-started birds and predator exclusion fencing. Their poor wild productivity is due to high nest predation and spring flood events, which together with their small population size and range remains a cause for concern.

Habitat grouping from LNRS priority list

Lowland fens, floodplain wet grasslands and associated habitats

Stone-curlew *Burhinus oedicnemus*

►► Actions

ACTION 1 ► expand breeding population of Stone-Curlew within East Cambridgeshire through:

- creating grasslands similar to the Breckland area, with grass kept no taller than 2 cm from March to July, so the birds can easily nest and spot predators.
- encouraging more rabbits, as their burrows help create a good nesting area for the birds.
- protecting the birds from predators by implementing predator control measures, including fencing to keep them away from nest sites.

ACTION 2 ► create two-hectare nesting areas within 5 km of breeding sites, kept mostly bare/ sparsely vegetated ground by disturbing the ground every spring.

SUPPORTING ACTION 1 ► collaborate with stakeholders (e.g. RSPB) to investigate how Stone Curlews use the East Cambridgeshire landscape,

to better understand it's importance to the Stone-curlew population of Breckland Special Area of Conservation.

SUPPORTING ACTION 2 ► work with landowners to encourage effective usage of agri-environment schemes to encourage update of options that promote stone curlew recovery.

Justification for inclusion

A rare bird nationally. East Cambridgeshire is on the edge of its stronghold in The Brecks, so it is rare in Cambridgeshire and Peterborough. Still under threat despite conservation efforts as vulnerable to disturbance and predation.

Habitat grouping from LNRS priority list

Farmland mosaic (all)



Stone-curlew

Swift *Apus apus*

►► Actions

ACTION 1 ► provide nesting spaces that are suitable for Common Swifts and House Martins and do not disturb, remove nests, or try to limit these birds from nesting.

ACTION 2 ► install integrated bird boxes onto all new commercial and community buildings and 25% of new dwellings (where appropriate), secured as part of planning applications. This should include swift boxes positioned under the eaves of a building at least 5 metre above the ground with easy access.

ACTION 3 ► create insect-rich habitats near nesting sites, as well as at a larger landscape-scale.



SUPPORTING ACTION ►

work with partners to raise awareness about the plight of swifts and importance of creating new nesting sites with members of the public and local communities.

Justification for inclusion

Significant population declines have occurred. A lack of nest sites is thought to be the key driver of decline. Decline in their insect food has also thought to have contributed.

Habitat grouping from LNRS priority list

Urban landscape mosaic

Turtle Dove *Streptopelia turtur*

►► Actions

ACTION 1 ► create and maintain tall scrubby hedgerows and/or dense scrub thickets for nesting, and provide freshwater sources (e.g. ponds, streams) in proximity to the newly created habitat

ACTION 2 ► create suitable habitat in farmland landscape by providing areas of uncropped field edges or setting aside plots of farmland.

ACTION 3 ► plant buffer strips around farm fields with wild bird seed mix or scatter seeds for birds.

ACTION 4 ► restore or create semi-natural grassland.

SUPPORTING ACTION 1 ► manage hedges to provide nesting opportunities and allow suitable areas of scrub to grow.

SUPPORTING ACTION 2 ► work with landowners to encourage effective usage of agri-environment

schemes to encourage update of options that promote turtle dove recovery.

Justification for inclusion

One of the UK's fastest declining birds. East of England is still a stronghold. Scrub is required for breeding success. Decline caused by habitat loss and a reduction in food availability from changes in agriculture and increases in herbicide use. This has resulted in a shorter breeding season ultimately causing fewer nesting attempts. Turtle doves are also vulnerable to hunting pressures on migration and in African wintering grounds, where they may be affected by land use change.



Habitat grouping from LNRS priority list

Farmland (all)


Fairy Shrimp *Chirocephalus diaphanus*

►► Actions

ACTION 1 ► allow for disturbance and maintain winter water levels (although eggs can remain viable in the soil for many years without water).

ACTION 2 ► creation of scrapes (shallow depressions or disturbed areas in the ground) close to recorded sites and in places where temporary ponds may form.

SUPPORTING ACTION 1 ► work with local partners and recorders to undertake surveys to confirm presence.



Justification for inclusion

Few recent records nationally and only one site known in Cambridgeshire. Most recent record for this site is one from an individual reared from a soil sample in 2008. So continued presence needs confirming but included on the understanding it may not yet be extinct at the site.

Habitat grouping from LNRS priority list

Ponds

White-clawed Crayfish *Austropotamobius pallipes*

►► Actions

ACTION 1 ► prioritise their habitat needs by achieving consistent, steady flows of good or very good quality water.

ACTION 2 ► manage riverbanks to offer numerous natural or artificial 'refuges' which offer opportunities to hide from predators.

ACTION 3 ► establish ark sites (protected areas) to maintain the number of populations, in accordance guidance (e.g. Crayfish Conservation Manual)

SUPPORTING ACTION 1 ► survey watercourses to confirm distribution and introduction of White Clawed Crayfish and inform landowners, developers and other stakeholder of their presence.

SUPPORTING ACTION 2 ► coordinate with



Neighbouring Authorities that have identified this species as part of their LNRS and work on including support development and research into a vaccine.

SUPPORTING ACTION 3 ► promote biosecurity (e.g. for fishing gear) to reduce the risk of spreading diseases such as crayfish plague.

Justification for inclusion

Globally endangered species and severely threatened in the UK. In Cambridgeshire and Peterborough small populations may still be present in a few isolated locations.

Habitat grouping from LNRS priority list

Chalk streams

Autumn Lady's Tresses *Spiranthes spiralis*

►► Actions

ACTION 1 ► manage sites through management practices such as grazing and cutting to maintain a short grassland towards the end of the growing season, allowing plants to flower and set seed.

SUPPORTING ACTION 1 ► reduce the usage of herbicide and fertiliser where possible by working with local landowners.



Autumn Lady's Tresses

Justification for inclusion

Only present at 1-2 sites in Cambridgeshire and Peterborough. Good indicator of unimproved grassland. Significant national decline.

Habitat grouping from LNRS priority list

Lowland calcareous grassland

Bastard-toadflax *Thesium humifusum*

►► Actions

ACTION 1 ► manage chalk and limestone grassland through management practices such as grazing and cutting to create a short grassland sward.

SUPPORTING ACTION 1 ► reduce the usage of herbicide and fertiliser where

possible, working with local landowners.

Justification for inclusion

Strong calcareous grassland indicator. Restricted distribution nationally. Present at 2-3 sites in Cambridgeshire and Peterborough. Requires short open grassland so always under threat from a lack of management and lower levels of grazing.

Habitat grouping from LNRS priority list

Lowland calcareous grassland



Bastard-toadflax

Broad-leaved Cudweed *Filago pyramidata*

►► Actions

ACTION 1 ► create regular disturbance to expose soil for seedlings to grow.

SUPPORTING ACTION 1 ► protect known sites against changes of nutrients in the soil (nutrient enrichment) by working with local landowners to reduce the usage of herbicide and fertiliser where possible.



Broad-leaved Cudweed

Justification for inclusion

GB and England Red List Endangered. Only two post 2000 sites in Cambridgeshire and Peterborough (last records 2012). Included on the understanding that it is still present.

Habitat grouping from LNRS priority list

Arable field margins


Cambridge Milk-parsley *Selinum carvifolia*

►► Actions

ACTION 1 ► work with local partners to monitor existing known locations of this species.

ACTION 2 ► ensure that scrub is managed well, and the water table is kept at optimum levels (close to the surface, but not so high that it submerges the plant completely).

SUPPORTING ACTION 1 ► undertake a feasibility assessment regarding whether there are suitable sites for the reintroduction of this species.



Cambridge Milk-parsley

Justification for inclusion

GB and England Red List Endangered. Cambridgeshire holds the only populations nationally. May now only be present at 1-2 native sites.

Habitat grouping from LNRS priority list

Lowland fens, floodplain wet grasslands and associated habitats

Chalk Fragrant orchid *Gymnadenia conopsea*

►► Actions

ACTION 1 ► ensure appropriate levels of cutting and/or grazing.

ACTION 2 ► promote the removal of scrub

Justification for inclusion

Strong chalk grassland indicator.

May only be present at two sites in Cambridgeshire and Peterborough now.

Habitat grouping from LNRS priority list

Lowland calcareous grassland.



Chalk Fragrant orchid


Common Butterwort *Pinguicula vulgaris*

►► Actions

ACTION 1 ► restore and / or maintain optimum water levels (just above the surface or level with the surface of the soil) to keep it moist without submerging the plant across the site.

ACTION 2 ► reduce and manage water pollution by partnering with landowners to improve practices on their land.

ACTION 3 ► control scrub taking over through management practices such as cutting and/or grazing.



Common Butterwort

Justification for inclusion

England Red List Vulnerable. At severe risk of extinction from Cambridgeshire and Peterborough at its last remaining native site if still present.

Part of a very rare habitat in our area at this site.

Habitat grouping from LNRS priority list

Lowland fens, floodplain wet grasslands and associated habitats



Common Cottongrass

Common Cottongrass *Eriophorum angustifolium*

►► Actions

ACTION 1 ► restore and / or maintain optimum water levels (typically 5cm – 10cm deep) across the site by working with stakeholders (e.g. Natural England).

ACTION 2 ► reduce and manage water pollution by partnering with landowners to improve practices on their land.

ACTION 3 ► control scrub taking over through appropriate levels of cutting and/or conservation grazing in spring and summer (taking care to prevent poaching during wet periods)

ACTION 4 ► consider the extension of existing sites

for recolonisation and the creation of new sites for translocation.

Justification for inclusion

England Red List Vulnerable. Only present at one remaining native site in Cambridgeshire and Peterborough as part of a very rare habitat in our area.

Habitat grouping from LNRS priority list

Lowland fens, floodplain wet grasslands and associated habitats

Crested Cow-wheat *Melampyrum cristatum*

►► Actions

ACTION 1 ► implement positive management and monitoring of sites where it is known to occur. Management includes cutting and removing the arisings at appropriate times of the year and buffer areas to prevent spray drift from adjacent farmland.

ACTION 2 ► protect road verges from damage / disturbance and work with Cambridgeshire Highways for implementation of better management.

ACTION 3 ► consider the extension of well-managed

road verges and new grassland areas to allow for recolonisation and the potential for translocation.

Justification for inclusion

GB and England Red List Endangered. Ancient woodland edge species, remaining on some verges. Southern Cambridgeshire and north-west Essex is the centre of its national distribution. At severe risk of extinction at most remaining sites.

Habitat grouping from LNRS priority list

Woodland (All)



Crested Cow-wheat

Dodder *Cuscuta epithymum*

►► Actions

ACTION 1 ► maintain chalk and limestone grassland by working with partners to ensure regular scrub management is achieved through mowing, cutting or grazing.

ACTION 2 ► prevent overgrazing by working with local landowners

SUPPORTING ACTION 1 ► reduce the usage of herbicide and fertiliser where possible by working with local landowners.

SUPPORTING ACTION 2 ► promote complementary

management practices by working with Rail and Highways Authorities.

Justification for inclusion

GB and England Red List Vulnerable. May now only be present at 1-2 sites in Cambridgeshire and Peterborough. An annual that is parasitic on the stems of a wide variety of small shrubs and herbs.

Habitat grouping from LNRS priority list

Grassland (All)



Dodder

Dandelion

English Dandelion *Taraxacum anglicum*

Hampshire Dandelion *Taraxacum akteum*

Marsh Dandelion *Taraxacum palustre*

►► Actions

ACTION 1 – work with local partners to monitor existing known sites.

ACTION 2 ► ensure positive management through suitable grazing and mowing.

ACTION 3 ► permitting winter flooding and monitoring water quality (or change to vegetation) to ensure eutrophication is countered by management.

Justification for inclusion

Taraxacum anglicum: confined to a small number of calcareous fens and water-meadows in central and southern England, with a high-water table and winter flooding, but probably intolerant of eutrophication

Taraxacum akteum: Confined to species-rich undisturbed meadows with a high-water table in approximately 10 sites in central England

Taraxacum palustre: confined to a small number of calcareous fens and water-meadows in central and

southern England, with a high-water table and seasonally inundated hay meadows, but probably intolerant of eutrophication. Also found in a few coastal grasslands and peaty grasslands. Main concentration of sites are Cambridgeshire and the New Forest.

Habitat grouping from LNRS priority list

Grassland and urban landscape (all)



Fen Wood-rush *Luzula pallescens*

►► Actions

ACTION 1 ► monitoring of sites where it is known to occur and positive management which includes disturbance of peaty areas at appropriate times.

ACTION 2 ► create suitable, additional areas of disturbance to allow for the recovery of the species.

Justification for inclusion

GB and England Red List Critically Endangered. Cambridgeshire holds the only populations nationally at Holme Fen and Woodwalton Fen. It is a short-lived species that requires disturbance so under threat from a lack of management, but has a persistent seed bank.

Habitat grouping from LNRS priority list

Lowland fens, floodplain wet grasslands and associated habitats



Fibrous Tussock-sedge *Carex appropinquata*

►► Actions

ACTION 1 ► positive management of site where it is known to occur, especially the specific areas where it is found, including sufficient levels of grazing and/or cutting and maintaining appropriate water levels.

ACTION 2 ► consider where the extension of cutting and grazing regimes could allow for the expansion of this species.



Fibrous Tussock-sedge

Justification for inclusion

Restricted in distribution nationally and only found at one site in Cambridgeshire and Peterborough (Wicken Fen). Evidence of a decline there in recent decades.

Habitat grouping from LNRS priority list

Lowland fens, floodplain wet grasslands and associated habitats


Flat-sedge *Blysmus compressus*

►► Actions

ACTION 1 ► maintain an open, short grassland through either an extensive grazing regime or by annual mowing and the removal of arisings. Periodic removal of scrub taking over may be necessary.

SUPPORTING ACTION 1 ► allow for the trampling of informal footpaths, as has a competitive advantage when soil is waterlogged or compacted.

SUPPORTING ACTION 2 ► efforts should be made to ensure a dynamic hydrological regime and the restoration of natural function where possible.



Flat-sedge

SUPPORTING ACTION 3

- reduce the usage of herbicide and fertiliser where possible by working with local landowners.

Justification for inclusion

GB and England Red List Vulnerable. At severe risk of extinction from Cambridgeshire and Peterborough as only now present in small numbers at its last remaining site.

Habitat grouping from LNRS priority list

Lowland fens, floodplain wet grasslands and associated habitats

Fly Orchid *Ophrys insectifera*

►► Actions

ACTION 1 ► where fly orchid is known / likely to be present, manage scrub along the edges, rides, and glades of woodland to create dappled light conditions and thin grassland conditions under a canopy of open scrub or grassland canopy.

Justification for inclusion

GB and England Red List Vulnerable. Only one remaining extant site in Cambridgeshire and Peterborough so under threat. Can be found in

a range of habitats but our population is within a woodland in the NW of the area. Formerly found in southern Cambridgeshire but now lost from all its former sites there.

Habitat grouping from LNRS priority list

Woodland (all)



Fly Orchid

Frog Orchid *Coeloglossum viride*

►► Actions

ACTION 1 ► maintain meadows and pasture by working with partnership (e.g. Natural England, Wildlife Trust) through management practices such as mowing, cutting and conservation grazing.

ACTION 2 ► restore and create suitable historic and new habitat sites.



Frog Orchid

Justification for inclusion

GB and England Red List Vulnerable. Only present at two sites so at risk of extinction in Cambridgeshire and Peterborough.

Habitat grouping from LNRS priority list

Lowland calcareous grassland

Great Pignut *Bunium bulbocastanum*

►► Actions

ACTION 1 ► maintain calcareous grassland by working with partners (e.g. Wildlife Trust) to ensure regular scrub management is achieved through mowing, cutting and grazing.

ACTION 2 ► restore and create suitable historic and new habitat sites

SUPPORTING ACTION 1 ► create some disturbance to expose soil for seedlings to grow but allow mature plants to thrive in areas with more cover.



Great Pignut

Justification for inclusion

Nationally restricted to a few sites in Hertfordshire, Buckinghamshire, Bedfordshire and Cambridgeshire.

Now may only be present at the Cherry Hinton Chalk Pits in Cambridgeshire and Peterborough.

Habitat grouping from LNRS priority list

Lowland calcareous grassland

Goldilocks Buttercup *Ranunculus auricomus* group

Bourn Goldilocks Buttercup

Ranunculus aequilaterus

Angular Goldilocks Buttercup

Ranunculus angularis

Boulder Clay Goldilocks Buttercup

Ranunculus argillicola

Buff Goldilocks Buttercup

Ranunculus armingfordensis

Backs Goldilocks Buttercup

Ranunculus cantabrigiensis

Comberton Goldilocks Buttercup

Ranunculus combertonensis

Carlton Goldilocks Buttercup

Ranunculus crassilobus

Henslow's Goldilocks Buttercup

Ranunculus henslowii

Hertford Goldilocks Buttercup

Ranunculus hertfordensis

Gransden Goldilocks Buttercup

Ranunculus obovatilobatus

Caxton End Goldilocks Buttercup

Ranunculus pubitorus

Madingley Goldilocks Buttercup

Ranunculus rotundilobus

Stalked Goldilocks Buttercup

Ranunculus stipitatus

Toft Goldilocks Buttercup

Ranunculus toftensis

Walters's Goldilocks Buttercup

Ranunculus waltersii

Goldilocks Buttercup



►► Actions

ACTION 1 ► cut and collect cuttings in late summer / autumn after the buttercups have flowered and formed seeds.

ACTION 2 ► promote management practices for the buttercups (where they are found) over amenity value or gardening for other species, such as avoid mowing or strimming before buttercups have finished flowering and forming seeds.

Justification for inclusion

For historical reasons, due to researchers based in Cambridgeshire, the microspecies goldilocks-buttercups, *Ranunculus auricomus* species, are better known in Cambridgeshire than elsewhere in Britain.

However, for different reasons, there is good

evidence that they merit conservation action, and that their Cambridgeshire bias in distribution is real, and not an artefact of recording.

None of the British species of goldilocks-buttercups have so far been recorded in mainland Europe, where many more small-range endemic species are described; so, they represent small-range endemics in a global context.

Of the 15 microspecies recorded from Cambridgeshire, 13 are known only from county, and one from Cambridgeshire and Hertfordshire. The species shown here appear to be rare, and to have very restricted distributions. Surveys in neighbouring counties have failed to find the Cambridgeshire endemic species.

Habitat grouping from LNRS priority list

Woodland (all), Grassland (all)

Heath Dog-violet *Viola canina* subsp. *ruppii*

►► Actions

ACTION 1 ► positive management and monitoring of the sites where it is known to occur, along with the restoration and creation of historical and new sites, which includes the appropriate levels of cutting and/or grazing and monitoring of water levels.

ACTION 2 ► create disturbance to encourage germination.

Justification for inclusion

GB and England Red List Endangered.



Heath Dog-violet

This sub-species of heath dog-violet grows in wetter habitats than sub-species *canina* and is found in only two sites nationally, both in Huntingdonshire (Holme Fen and Woodwalton Fen). It is therefore under threat.

Habitat grouping from LNRS priority list

Lowland fens, floodplain wet grasslands and associated habitats

Maiden Pink *Dianthus deltoides*

►► Actions

ACTION 1 ► restore and create suitable historic and new habitat sites, gravel-topped chalk hills or gravel extracted pits are preferred.

ACTION 2 ► ensure appropriate management practices at sites, with light grazing and occasional disturbance to promote the growth of new plants from seeds.

SUPPORTING ACTION 1 ► work with local partners to monitor known and new sites where it is known to occur.

Justification for inclusion

England Red List Vulnerable. Still present but under threat at only remaining native site at Hildersham in Cambridgeshire and Peterborough.

Habitat grouping from LNRS priority list

Lowland calcareous grassland



Maiden Pink

Marsh Fragrant-Orchid *Gymnadenia densiflora*

►► Actions

ACTION 1 ► implement positive management of wet grassland, fen, marsh and swamp sites where it is known to occur, along with the restoration and creation of historical and new sites, to encourage the right conditions, including appropriate levels of cutting and/ or grazing.


- monitoring and maintaining appropriate water levels
- control invasive plants that might outcompete marsh orchid for light, space and nutrients.
- preserve soil conditions

Justification for inclusion

Strong indicator of high-quality wet fen meadow habitats. At severe risk of extinction from Cambridgeshire and Peterborough as only present now at one or two sites.

Habitat grouping from LNRS priority list

Lowland fens, floodplain wet grasslands and associated habitats



Marsh Fragrant-Orchid

Marsh Helleborine *Epipactis palustris*

Actions

ACTION 1 ► implement positive management of site where it is known to occur, along with the restoration and creation of historical and new sites, which includes the appropriate levels of and monitoring of water levels to create flushed or seasonally inundated areas.

- appropriate levels of cutting and/or grazing
- monitoring and maintaining water levels to create flush or seasonally inundated areas.
- preserve soil conditions to be damp and nutrient poor (preferably peaty soils flushed



with calcareous groundwater)

- create and maintain open space with sparse vegetation.

Justification for inclusion

Strong indicator of high-quality wet fen meadow habitats. At severe risk of extinction from Cambridgeshire and Peterborough as only present now at one site.

Habitat grouping from LNRS priority list

Lowland fens, floodplain wet grasslands and associated habitats

Marsh Lousewort *Pedicularis palustris*

►► Actions

ACTION 1 ► create and enhance suitable areas to increase populations of Marsh Lousewort by

- maintain moist, waterlogged conditions where soil remains consistently wet, during growing season (early spring and extends through summer)
- create shallow, boggy areas with nutrient poor acidic peaty soils to promote plants growth.
- control invasive vegetation that might outcompete Marsh Lousewort for light, space and nutrients. By appropriate levels of cutting/ or grazing.
- encourage natural water flow, with seasonal fluctuations.



SUPPORTING ACTION 1 ► Manage water quality by working with landowners to reduce the usage of herbicide and fertiliser where possible.

Justification for inclusion

England Red List Vulnerable. Only now present at likely two sites in Cambridgeshire and Peterborough. Would have once been relatively common in fens and bogs in Cambridgeshire and Peterborough. It has been lost from both Wicken and Chippenham Fens for example.

Habitat grouping from LNRS priority list

Lowland fens, floodplain wet grasslands and associated habitats

Milk-parsley *Thysselinum palustre*

►► Actions

ACTION 1 ► implement positive management and monitoring of the wet grassland, fen, marsh, swamp site where it is known to occur, along with the restoration and creation of historical and new sites by

- maintain appropriate levels of cutting (four-year sedge-cutting cycle) and/or grazing and monitoring of water levels.
- monitor habitat disturbances, by minimizing human activity.

Justification for inclusion

GB and England Red List Vulnerable. Fenland species, now only present in Cambridgeshire and Peterborough as a native at Wicken Fen but is not abundant there. Foodplant of swallowtail butterfly which requires substantial populations and large individual plants for its survival.

Habitat grouping from LNRS priority list

Lowland fens, floodplain wet grasslands and associated habitats



Milk-parsley



Moon Carrot

Moon Carrot *Seseli libanotis*

►► Actions

ACTION 1 ► ensure appropriate levels of cutting and/or grazing.

ACTION 2 ► promote the removal scrub.

ACTION 3 – promote practices to control rabbits.

SUPPORTING ACTION 1 ► work local partners to monitor calcareous grassland sites where it is known to occur.

Justification for inclusion

Very rare nationally, this species is only present in Cambridgeshire, Bedfordshire and Sussex. The Cambridgeshire populations are relatively small and are all centred in an area on the SE side of Cambridge. Some of these populations have declined in recent years.

Habitat grouping from LNRS priority list

Lowland calcareous grassland

Mountain Everlasting *Antennaria dioica*

Actions

ACTION 1 ► maintain a short, open sward, best achieved by sheep grazing in later summer and early autumn, allowing it and other associate species to flower and set seed in the spring and summer months.

SUPPORTING ACTION 1

► protect against damage from visitors and dog walkers using the nature reserve where it is known to occur.

SUPPORTING ACTION 2 ► look for opportunities to extend the species range.

SUPPORTING ACTION 3 ► seek out opportunities for creating calcareous grassland in John Clare



Countryside and particularly as chalk quarries are restored.

SUPPORTING ACTION 4

► include Mountain Everlasting in the restoration and management of the species assemblage.

Justification for inclusion

England Red List Vulnerable. Only found at one site in Cambridgeshire and Peterborough and rare in lowland England so level of threat high.

Habitat grouping from LNRS priority list

Lowland calcareous grassland

Pasqueflower *Pulsatilla vulgaris*

►► Actions

ACTION 1 ► maintain chalk and limestone grassland to an approximate height of 5cm and ensure a clear surrounding of vegetation.

ACTION 2 ► sites managed through conservation grazing (sheep or mixed livestock) by working with landowners and partners (e.g. Natural England). However, mixed livestock should not be present during flowering period to prevent overgrazing.



SUPPORTING ACTION 1 ► undertake further research on the colonisation of new sites and their dispersal mechanism.

Justification for inclusion

GB and England Red List Vulnerable. Strong calcareous grassland indicator. Only present now in Cambridgeshire and Peterborough at 2-3 native sites.

Habitat grouping from LNRS priority list

Lowland calcareous grassland.

Rare Spring-sedge *Carex ericetorum*

►► Actions

ACTION 1 ► maintain chalk and limestone grassland sites, where Rare Spring-sedge has been recently and historically known to occur, with appropriate levels of grazing to create open conditions and prevent scrub taking over.

ACTION 2 ► encourage grazing during the late summer, autumn and winter months, with the aim of producing short grassland for the spring and summer months, allowing flowering and seed formation for a wide range of species throughout the growing season.



Rare Spring-sedge

SUPPORTING

ACTION 1 ► avoid using fertilisers, manures, or spot treatments for weeds

Justification for inclusion

GB and England Red List Vulnerable. Only recent records from two sites in Cambridgeshire and Peterborough. Under threat from reductions in grazing levels.

Habitat grouping from LNRS priority list

Lowland calcareous grassland

Red-tipped Cudweed *Filago lutescens*

►► Actions

ACTION 1 ► create some disturbance in early autumn to expose soil for seedlings to grow.

ACTION 2 ► reduce the fragmentation of suitable habitats and extend range of pre-existing farmed landscape habitats managed for arable plants.

SUPPORTING

ACTION 1 ► continue to work with local partners to monitor and understand the Red-Tipped Cudweed status at the last



Red-tipped Cudweed

known site in Cambridgeshire and Peterborough.

Justification for inclusion

GB and England Red List Endangered. Only one post 2000 site for this species (last record 2004) so may now be extinct in Cambridgeshire and Peterborough. Included on the understanding that it still may be present.

Habitat grouping from LNRS priority list

Arable field margins

Slender Hare's-ear *Bupleurum tenuissimum*

►► Actions

ACTION 1 ► positive management of saltmarsh habitat along the River Nene north of Wisbech, including appropriate management practices such as cutting and/or grazing to reduce the dominance of couch grass species.

Justification for inclusion

GB and England Red List Vulnerable. May now only be present in Cambridgeshire and Peterborough

in the restricted habitat of saltmarsh/coastal grassland north of Wisbech by the River Nene and under threat there from being swamped out by coarse grasses due to a lack of grazing.

Habitat grouping from LNRS priority list

Lowland calcareous grassland



Slender Hare's-ear

Smooth Rupturewort *Herniaria glabra*

►► Actions

ACTION 1 ► positive management of native site where it has been recently known to occur, which includes disturbance and creation of open ground.

Justification for inclusion

Native national distribution is restricted mostly to the Breckland area. Thought to likely still be present



Smooth Rupturewort

at one remaining native location in Cambridgeshire (gravel pits near Kennett). Can be found elsewhere as a garden escape or accidental introduction. Management for this species may help other Breckland/sandy ground rarities.

Habitat grouping from LNRS priority list

Grassland (all)

Spanish Catchfly *Silene otites*

►► Actions

ACTION 1 ► ensure appropriate management practices on sites where it is known to occur, with appropriate levels of cutting and/or conservation grazing and occasional disturbance to promote the growth of new plants from seeds.

ACTION 2 ► work with partners (e.g. Wildlife Trust) to monitor site and identify optimal management for this species.

ACTION 3 ► create any new sites within close proximity to existing populations.

Justification for inclusion

GB and England Red List Endangered. National distribution is restricted to the Breckland area. Now thought to be lost from its native locations at the eastern edge of Cambridgeshire but still present in a small area on an old railway line near Burwell.

Here it is thought to be an accidental introduction with ballast imported from the Freckenham area where it occurred naturally. This small, isolated population is under threat.

Habitat grouping from LNRS priority list

Lowland calcareous grassland



Spanish Catchfly

Spotted Cat's-ear *Hypochaeris maculata*

►► Actions

ACTION 1 ► ensure appropriate management of calcareous grassland, with conservation grazing and scrub management, to create an open area. Work with partners (e.g. Natural England) to monitor site and identify optimal management for this species.

ACTION 2 ► restore and create additional grassland habitat.

ACTION 3 ► protect from over grazing at certain times of the year and control scrub.

Justification for inclusion

GB and England Red List Vulnerable. Strong chalk grassland indicator. May only be present at one site in Cambridgeshire and Peterborough with a small population.

Habitat grouping from LNRS priority list

Lowland calcareous grassland



Spotted Cat's-ear

Violet Helleborine *Epipactis purpurata*

►► Actions

ACTION 1 ► provide dense shade and, when felling, areas adjacent to populations are left in dense shade also. Refrain from creating any new areas of glade in areas where populations of violet helleborine are known to occur.


ACTION 2 ► protection of flowering spikes from grazing by rabbits and deer.

Justification for inclusion

Only present at 2-3 sites in Cambridgeshire and Peterborough, in relatively small numbers so under severe threat.

Habitat grouping from LNRS priority list

Lowland mixed deciduous woodland



Violet Helleborine

Black Hairstreak Butterfly *Satyrion pruni*

Actions

ACTION 1 ► retain existing thickets of blackthorn and mature hedges with blackthorn.

ACTION 2 ► plant new thickets and hedgerows containing blackthorn.

ACTION 3 ► carry out management and coppicing routines for hedges, trees, and thickets which are suitable to the species.

ACTION 4 ► increase the connectivity of suitable habitats by creating and extending stands, trees, and hedgerows containing blackthorn which connect existing areas.

ACTION 5 ► create wide rides, glades, and scrub edges in and around woodlands.

ACTION 6 ► ensure maintenance of suitable nectar



Black Hairstreak Butterfly

sources, especially bramble microspecies which flower during the flight period.

Justification for inclusion

GB Red List Endangered. Significant decline and national distribution is restricted to a small belt of central England running approximately from Oxford to Peterborough, so populations in Cambridgeshire and Peterborough are nationally significant. Habitat is mature blackthorn scrub growing in sunny, sheltered areas.

Habitat grouping from LNRS priority list

Woodland (all)

Caddis fly *Eretsis baltica*

Actions

ACTION 1 ► maintain constant water levels at sites where this caddisfly is known to occur ensuring there is sufficient aquatic vegetation and roots of marginal trees.

ACTION 2 ► dig small pools in dry summers to provide temporary refuges.

ACTION 3 ► expand populations through further habitat creation.

SUPPORTING ACTION 1 ► See King's Dyke Nature Reserve for best practice examples of where they are creating clean water ponds at a restored mineral extraction site for freshwater wildlife.



Caddis fly

Justification for inclusion

GB Red List Vulnerable. Caddisfly whose larvae need permanently flooded fen and similar habitats. They are found amongst the submerged parts of dense emergent vegetation. Recorded in Cambridgeshire and Peterborough from two sites, although has not been recorded at one (Wicken Fen) for several decades.

Habitat grouping from LNRS priority list

Lowland fens, floodplain wet grasslands and associated habitats



Desmoulin's Whorl Snail

Desmoulin's Whorl Snail *Vertigo (Vertigo) moulinsiana*

►► Actions

ACTION 1 ► maintain water levels on wetland sites at water-table or slightly above ground surface for much of the year.

ACTION 2 ► ensure the wetted zone does not become over-shaded or scrubbed up.

ACTION 3 ► manage vegetation through appropriate management practices of cutting and/or grazing.

SUPPORTING ACTION 1 ► in river systems, focus on protecting and supporting populations upstream first.

Justification for inclusion

GB Red List Vulnerable. National decline and recent records only in 1-2 locations in Cambridgeshire and Peterborough but has been found to be under recorded nationally. It is restricted to calcareous wetlands, usually bordering lakes or rivers, or in fens and so is a positive indicator species for this type of habitat.

Habitat grouping from LNRS priority list

Lowland fens, floodplain wet grasslands and associated habitats

Eyed Longhorn Beetle *Oberea oculata*

►► Actions

ACTION 1 ► create, restore and enhance wet woodland with a particular focus on maintenance of suitable willow habitat.

ACTION 2 ► ensure a variety of habitats, including dead wood, healthy trees, young saplings, scrub, and open spaces. Keep all dead wood (standing and fallen) where it is and preserve trees with decay features without damaging them.

ACTION 3 ► buffer and connect fragmented woodland using suitable tree species.

ACTION 4 ► remove invasive and competitive plant species known to negatively impact wet woodland habitat such as giant hogweed and Japanese knotweed.

SUPPORTING ACTION 1 ► support actions to locate, develop and implement a species recovery plan to build a resilient population, working with partners such as Buglife.



Eyed Longhorn Beetle

Justification for inclusion

GB Red List Critically Endangered. Cambridgeshire previously thought to hold the only site nationally for this species (there also appears to be some recent confirmed iRecord records for a site in London).

Its requirements are young/regenerating *Salix* so possible sites can be found and/or created in gravel pits and other new wetlands. A possible flagship species for the fauna of invasive willows/osier beds, with implications for gravel pit restoration and habitat creation.

Habitat grouping from LNRS priority list

Lowland fens, floodplain wet grasslands and associated habitats


Ground Beetle *Ophonus puncticollis*

►► Actions

ACTION 1 ► support moss habitats by managing with cutting, grazing, and scrub control. Also, create some disturbance to keep areas with mostly annual and biennial plants.

Justification for inclusion

GB Red List
Endangered. Very rare nationally. The last Cambridgeshire record was in 1989 from Devil's Dyke and it may still occur there. The other rare chalk grassland *Ophonus*, of which Cambridgeshire and Peterborough used to have several, have already



Ground Beetle

largely disappeared and it could be argued that this is one of the last representatives of that fauna. This reflects the decline in extent and quality of chalk grassland habitat in Cambridgeshire and Peterborough. Succession is a threat and management for this species would benefit other species even if not still present.

Habitat grouping from LNRS priority list

Lowland calcareous grassland.

Large-mouthed Valve Snail *Valvata macrostoma*

►► Actions

ACTION 1 ► positive management of drains and waterways where it is known to exist, which includes avoiding eutrophication (particularly nitrate / nitrite) and drain management at appropriate times and intervals.

SUPPORTING ACTION 1 ► undertake surveys to look for other potential existing populations, establishing its status in the county, along with suitable sites to establish new populations.

Justification for inclusion

GB Red List Vulnerable. National distribution is scattered and local, and it is thought there has been a significant decline. In Cambridgeshire and Peterborough recent records for the Ouse and Nene Washes and 1990s Wicken Fen records. Found in vegetated drains with relatively good water quality so under threat from eutrophication.

Habitat grouping from LNRS priority list

Fen drainage ditches



Large-mouthed Valve Snail

Marsh Carpet Moth *Gagitodes sagittata*

►► Actions

ACTION 1 ► implement the following positive management of the wetland sites where it is known to occur, along with the restoration and creation of historical and new sites to encourage the right conditions for common meadow-rue.

- maintain wet, undisturbed conditions.
- control water levels, avoiding excessive flooding.
- control invasive plants that might outcompete the meadow-rue.
- avoid nutrient runoff and pollution.

Justification for inclusion

Restricted distribution nationally with most records in East Anglia. Found in Cambridgeshire and Peterborough at Wicken Fen and some other wetland localities. Larval food plant meadow-rue is relatively rare so it is restricted because of this.

Habitat grouping from LNRS priority list

Lowland fens, floodplain wet grasslands and associated habitats

Marsh Carpet Moth

Mud Beetle *Sphaerius acaroides*

►► Actions

ACTION 1 ► implement positive management of the site where it is known to occur. The species needs fluctuating water levels on mid-successional stages with bare/alga coated mud, so rotational management is needed.

Justification for inclusion

GB Red List Endangered. A small beetle found around the margins of waterbodies. Very rare nationally but may be overlooked. Only one site in Cambridgeshire and Peterborough, Orton Pits. This site is threatened by successional change. The old Peterborough brick pits are nationally important for a range of taxonomic groups and this could be a flagship species as management for this beetle would also benefit other species.

Habitat grouping from LNRS priority list

Lowland fens, floodplain wet grasslands and associated habitats


Pardosa Paludicola Spider

Pardosa paludicola

►► Actions

ACTION 1 ► implement positive management of the fen habitat where it is known to occur and surrounding sites, which includes the appropriate levels of cutting and/or grazing, and monitoring and maintaining a high-water table.

ACTION 2 ► reduce the fragmentation and extend the range of suitable habitats.



Pardosa Paludicola Spider

Justification for inclusion

GB Red List Endangered. Very rare nationally and only found at Woodwalton Fen in Cambridgeshire and Peterborough. A wolf spider that is found in damp habitats.

Habitat grouping from LNRS priority list

Lowland fens, floodplain wet grasslands and associated habitats

Tansy Beetle *Chrysolina graminis*

►► Actions

ACTION 1 ► manage mosaic habitat favourable to the Tansy plant through grazing (being careful to avoid overgrazing).

ACTION 2 ► tackle invasive non-native species, such as Himalayan balsam, by working with Natural England, Buglife and other partners.

ACTION 3 ► seek to establish a population in suitable vegetation in the Great Fen on land which is not subject to flooding.

SUPPORTING ACTION 1 ► summer flooding should be managed as a threat to populations and working with catchment land managers is encouraged.



Tansy Beetle

Justification for inclusion

GB Red List Endangered. Very few native sites known nationally with the largest population along a 45km stretch of the River Ouse in central Yorkshire. Rediscovered in Cambridgeshire at Woodwalton Fen in 2014. Already subject to a species recovery and introduction programme coordinated by the Species Recovery Trust.

Habitat grouping from LNRS priority list

Lowland fens, floodplain wet grasslands and associated habitats



Taylor's Spire Snail *Marstoniopsis insubrica*

►► Actions

ACTION 1 ► positive management of drains and waterways where it has been previously recorded, which includes avoiding eutrophication and drain management at appropriate levels, times and intervals.

SUPPORTING ACTION 1 ► undertake new surveys to establish status.

Justification for inclusion

GB Red List Endangered. Nationally only recorded recently from a small number of locations. In Cambridgeshire and Peterborough last recorded in the Little Ouse river, but has been also recorded in the Great Ouse.

Found in slow moving water in rivers and canals in aquatic weeds. Under threat from habitat degradation.

Habitat grouping from LNRS priority list

Rivers; Fen main rivers and drains

Witham Orb Mussel *Sphaerium solidum*

►► Actions


ACTION 1 ► implement positive management of drains and waterways where it has been previously recorded, which includes avoiding eutrophication and drain management at appropriate levels, times and intervals.

SUPPORTING ACTION 1 ► undertake new surveys to establish status.

SUPPORTING ACTION 2 ► control the spread of the invasive species Asian Clam (*Corbicula fluminea*).

Justification for inclusion

GB Red List Critically Endangered. Nationally post-2000 records only in the fenland area. Recorded



at several sites in the New Bedford River and River Great Ouse in the early 2000s and last record in 2010. Surveys in 2015 at the same sites failed to find any live specimens so possibly now extinct. Further work, including eDNA surveys could help confirm status.

Threats are from eutrophication, saline intrusion, inappropriate drain and river management and the invasive mollusc *Corbicula fluminea*.

Habitat grouping from LNRS priority list

Rivers; Fen main rivers and drains

Zora Armillata Spider *Zora armillata*

►► Actions

ACTION 1 ► manage fen habitat through the appropriate levels of cutting and/or grazing. Work with partners (e.g. National Trust) to monitor site and identify optimal management for this species.

ACTION 2 ► monitor and maintain water levels

Justification for inclusion

GB Red List Critically Endangered. Very rare nationally - has previously been found in wet heath and bog in Dorset and in fen habitat in East Anglia.

Only found at Wicken Fen in Cambridgeshire and Peterborough.

Habitat grouping from LNRS priority list

Lowland fens, floodplain wet grasslands and associated habitats



Zora Armillata Spider

Hazel Dormouse *Muscardinus avellanarius*

►► Actions

ACTION 1 ► connect suitable habitat within 2 km of these sites by creating suitable connecting habitats e.g. coppice, woodland, or thick hedgerows.

ACTION 2 ► ensure arboreal connections across woodland rides every 50-100m and erect dormouse boxes and/or tubes.

SUPPORTING ACTION 1 ► retain trees which have cracks, crevices, and deadwood and create sunny areas in woodlands and retain woody species like blackthorn and hazel.

SUPPORTING ACTION 2 ► do not clear understorey in winter and do not clear fell in dormouse locations.

Justification for inclusion

GB Red List Vulnerable. Historic populations of this species in Cambridgeshire and Peterborough uncertain. Now only exists as an introduced species at a few sites. Included at the suggestion that more can be done to improve the size and sustainability of these populations..

Habitat grouping from LNRS priority list

Lowland mixed deciduous woodland and hedgerows



Hazel Dormouse

Adder *Vipera berus*

►► Actions

ACTION 1 ► create new areas that include scrub, rough grassland and deciduous woodland habitat or manage and improve existing mosaic habitats to allow the natural movement of adders.

ACTION 2 ► manage these areas to create different shapes and features, like paths and clearings. Vary the ground to provide sunny spots for basking and places for hibernation.

ACTION 3 ► improve habitat connectivity and encourage the creation of additional populations in the vicinity of existing, isolated populations and link with neighbouring nature reserves.

SUPPORTING ACTION 1 ► limit the presence of game birds, house cats and human disturbance (e.g. from dog walkers) within the critical parts of



Adder

these mosaic habitats (i.e. near the known hibernation sites, scrub or bracken management can redirect footpaths to avoid sensitive areas). By use signage to inform the public of adder presence and reminding them to keep to existing paths to increase public awareness and appreciation and reduce disturbance

Justification for inclusion

Only one recent confirmed site for adder in Cambridgeshire and Peterborough so under threat.

Habitat grouping from LNRS priority list

Lowland calcareous grassland

Lichen *Bellicidia incompta*

►► Actions

ACTION 1 ► implement good woodland management practices creating well-lit areas and protect from felling.

ACTION 2 ► look for opportunities to reconnect existing populations through tree planting and hedgerow creation.

SUPPORTING ACTION 1 ► conserve and maintain the Elm trees and suckers on which this lichen depends.



Lichen

this species in Cambridgeshire and Peterborough. The strongest known colony is in Savages Spinney, where it grows on pole-size elm stems (sucker regrowth which has reached at least several inches in diameter).

Some microspecies of elm appear to be particularly prone to producing exudate from slight wounds and this is the case at Savages Spinney.

Justification for inclusion

GB Red List Vulnerable. Lichen which occurs on wound tracks (sap runs) of trees, and was most often found on elm trees prior to the arrival of Dutch Elm Disease. It has declined significantly since this time with the loss of most of these trees.

Now rare nationally and only a few known sites for

Lichens are an excellent indicator of good quality; clean air so measures taken to support this species are also measures which can improve air quality for people and the wider environment (e.g. reducing local air pollution levels).

Habitat grouping from LNRS priority list

Woodland (all)

Elm *Ulmus* spp

Cambridgeshire only:

Hayley Elm *Ulmus crenata*

Madingley Elm *Ulmus madingleyensis*

Bassingbourn Elm *Ulmus platyphylla*

Leathery-leaved Elm *Ulmus coriaceifolia*

Long-toothed Elm *Ulmus longidentata*

(found in Cambridgeshire and global
population fewer than 100 trees)

Long-toothed Elm *Ulmus longidens*

Fat-toothed Elm *Ulmus obesidens*

Hatley Elm *Ulmus sylvatica*

Bonhunt Elm *Ulmus acutissima*

Prominent-toothed Elm

Ulmus prominentidens

Dark-leaved Elm *Ulmus atrovirens*

Large-toothed Elm *Ulmus prionopylla*

Cut-leaved Elm *Ulmus incisa*

►► Actions

ACTION 1 ► conserve and manage hedgerows and fen droves

ACTION 2 ► establish elm nurseries for the most restricted and vulnerable Cambridgeshire microspecies.

Justification for inclusion

Elmwoods, elms lining fen droves, and roadside and hedgerow elms, remain more significant features of Cambridgeshire (and parts of Essex, Suffolk and Norfolk) than elsewhere in Britain. The diversity of smooth-leaved elms is far greater here than elsewhere (despite some detailed surveys elsewhere).

Many smooth-leaved elm microspecies have very restricted geographic distributions, and some of these, as listed, are confined to Cambridgeshire, or better represented here than anywhere else.

Currently 40 elm microspecies are considered native to Cambridgeshire, 5 of which are currently known only from the county, and 10 of which have a known population of fewer than 100 trees: all of these appear to be British endemics, with no records from mainland Europe.

The county also has some of the very few surviving veteran elms and elm pollards in Britain.

Habitat grouping from LNRS priority list

Lowland mixed deciduous woodland



Hayley Elm

Bearded Stonewort *Chara canescens*

►► Actions

ACTION 1 ► working with multiple different partners (e.g. water companies, farmers/ landowners) to reduce water pollution caused by sewage or fertiliser.

ACTION 2 ► control change in habitat by removing competing vegetation or return parts of ponds to suitable habitats for Bearded Stonewort by clearance or by temporarily lowering water table.

ACTION 3 ► creation of new ponds close to existing habitats to encourage early colonisers and movement of Bearded Stonewort

Justification for inclusion

GB Red List Endangered. Nationally found in some locations near the coast and its only inland sites are some of the old Peterborough's Oxford Clay brick pits. It is usually found in clear brackish water, but it is tolerant of a range of salinities.

Habitat grouping from LNRS priority list



Bearded Stonewort

Ponds

Dwarf Stonewort *Nitella tenuissima*

►► Actions

ACTION 1 ► manage open water habitats by working with partners to improve water quality by reducing water pollution caused by the addition of sewage or fertiliser.

ACTION 2 ► control competing vegetation with grazing or periodic vegetation clearance.


ACTION 3 ► create new ponds close by to provide new habitat for early colonisers.

ACTION 4 ► management should also be undertaken at suitable historic sites with the aim of regenerating this species from the spore-bank.

Justification for inclusion

GB Red List Endangered. Only recorded at Wicken Fen in Cambridgeshire and Peterborough. A plant of calcareous fenland where it occurs in shallow peaty pools and ditches. It requires bare peat surfaces and at Wicken Fen it has only appeared in the summer or autumn following peat cutting activities.

Habitat grouping from LNRS priority list



Dwarf Stonewort

Ponds

Tassel Stonewort *Tolypella intricata*

►► Actions

ACTION 1 ► reinstate management at existent sites.

ACTION 2 ► creation and maintenance of new sites through periodic clearance of ditches, scrub control and grazing.

ACTION 3 ► monitor and maintain water levels (few centimetres to about 1 metre from the surface)


SUPPORTING ACTION 1 ► work with local partners to complete pond survey in Cambridgeshire.

Justification for inclusion

GB Red List Endangered. Recent records at only one site in Cambridgeshire and Peterborough. A species of ephemeral pools and other waterbodies with alkaline water that dry out in summer. A winter annual which usually germinates in the autumn.

Habitat grouping from LNRS priority list

Ponds



Tassel Stonewort

Did you know?

Peterborough brick pits hold the main population in Britain of Bearded Stonewort (*Chara canescens*)




Corncrake *Crex crex*

►► Actions

ACTION 1 ► creation of grassland habitat managed with tall vegetation during summer and autumn and delaying mid-summer mowing of fields to allow Corncrakes to successfully nest and rear their chicks.

SUPPORTING ACTION 1 ► expand Corncrake populations at Nene Washes and Ouse Washes through continued support of the existing national captive breeding release programme.



Corncrake

Justification for inclusion

The massive decline of the UK population, and extinction in England, justified the introduction attempts at the Nene and Ouse Washes.

The fate of these should be assessed before more widespread reintroductions are attempted.

Habitat grouping

Farmland (all)

Fen Ragwort *Jacobaea paludosa*

►► Actions

ACTION 1 ► adopt nature-based solutions which allow for periodic flooding and return natural function to rivers and floodplains to assist efforts to expand current populations.

ACTION 2 ► preserve existing habitat and identify new sites with optimal habitat requirements (wet, waterlogged conditions, seasonally wet areas around waterbodies, open unshaded areas and nutrient rich soil)

ACTION 3 ► protect from deer and slugs to ensure it is not outcompeted by surrounding vegetation.

SUPPORTING ACTION 1 ► partner with Cambridgeshire County Council, Natural England and other landowners to look for opportunities to work together to ensure species expansion.



Fen Ragwort

Justification for inclusion

GB and England Red List Critically Endangered.

Only one native extant population nationally and this is in Cambridgeshire (a roadside ditch SSSI near Ely). Has been introduced to many sites but most introductions have failed.

Recent studies have shown greater seed viability and more opportunity for captive rearing and release.

Habitat grouping

Lowland fens, floodplain wet grasslands and associated habitats

Fen Violet *Viola stagnina*

►► Actions

ACTION 1 ► ensure limited competition from other plants.

ACTION 2 ► periodically disturb the soil to stimulate germination e.g. from livestock and wet soil is needed in winter but waterlogging in spring and summer is detrimental.

ACTION 3 ► establish functioning populations at different locations within existing fen nature reserves, and potentially in restored fen habitats.

Justification for inclusion

GB and England Red List Critically Endangered.



Fen Violet

Very rare nationally. Found at only two native sites in Cambridgeshire and Peterborough and introduced to another. It has a sporadic occurrence and can reappear from the seed bank after many years dormant.

Raising in cultivation is very successful, so native seed sources are available for future reintroduction.

Habitat grouping

Lowland fens, floodplain wet grasslands and associated habitats

Large Marsh Grasshopper *Stethophyma grossum*

►► Actions

ACTION 1 ► re-introduce Large Marsh Grasshopper to historic range across Cambridgeshire fens (Great Fen, after which former sites at Wicken and Chippenham Fens).

Justification for inclusion

Currently classed as Near Threatened following an 85% reduction in its British range. The species currently survives in bogs and wet heaths in Dorset, Hampshire and Somerset. All these populations are vulnerable



Large Marsh Grasshopper

to climate change. Unless it is re-established in its former native range in East Anglia it is likely to become extinct in Britain.

Captive breeding has been successful, and self-sustaining reintroduced populations have been re-established at sites in Norfolk

Habitat grouping

Wetland (all)

Water Germander *Teucrium scordium*

►► Actions

ACTION 1 ► implement positive management and monitoring of the sites where it is known to occur, which includes appropriate levels of grazing and/or cutting and scrub control. Also monitoring of water levels and some disturbance.

Justification for inclusion

GB and England Red List Endangered. Very rare nationally, by the 1990s it was confined to three sites



Water Germander

in England - two in Cambridgeshire and one in Devon. Cambridgeshire sites under threat from a lack of management, scrub encroachment and shading. Introduced to Kingfisher's Bridge (next to one of the native sites) and this has been successful.

Habitat grouping

Lowland fens, floodplain wet grasslands and associated habitats

Pool Frog *Pelophylax lessonae*

►► Actions

ACTION 1 ► re-introduce Pool Frog population into its historic range in Cambridgeshire fens through identification and management of suitable sites, working with the Natural England and Amphibian and Reptile Conservation Trust national reintroduction project currently operating in Norfolk.

Justification for inclusion

The last native population, in Norfolk, became extinct in 1880s. It has since been re-established at its last Norfolk site. Cambridgeshire is considered to have been a significant part of its former native range, and suitable habitats are present.

Habitat grouping

Lowland fens, floodplain wet grasslands and associated habitats



Pool Frog

Large Copper *Lycaena dispar*

►► Actions

ACTION 1 ► Liaise with organisations such as Natural England, working on reintroduction programmes and learn what success is and bespoke habitat requirements for this species that could potentially be replicated in Cambridgeshire in the long-term.

Justification for inclusion

The species, in the form of an endemic subspecies, became extinct



Large Copper

in Britain in the 1860s. Repeated attempts to re-establish it in the Cambridgeshire Fens have so far failed. But a better understanding of its ecology may make a future attempt feasible if suitable habitat, extensive foodplant populations, and water level management is in place.

Habitat grouping

Lowland fens, floodplain wet grasslands and associated habitats

Swallowtail Butterfly *Papilio machaon*

►► Actions

ACTION 1 ► Liaise with organisations working on reintroduction programmes and learn what success is and bespoke habitat requirements for this species that could potentially be replicated in Cambridgeshire in the long-term.


ACTION 2 ► Work towards creating appropriate, large scale wetland habitats, to support sufficient, established food sources and enable future natural colonisation.

Justification for inclusion

The British Swallowtail is regarded as an endemic subspecies, confined to fenland habitats and Milk-parsley is the larval food plant. It is current confined to the Norfolk Broads but occurred in Cambridgeshire until the 1950s. Previous reintroductions to Cambridgeshire have failed through being unable to sustain sufficiently large populations of the larval food plant.

Habitat grouping

Lowland fens, floodplain wet grasslands and associated habitats



Swallowtail Butterfly

Fen Orchid *Liparis loeselii*

►► Actions

ACTION 1 ► re-introduce Fen Orchid population through restoration of fen and marsh habitats on sites where Fen Orchid was previously known. Provide wet areas with bare sand, short grasses and calcium-rich habitat upon which the Fen Orchid depend, working with partners such as Plantlife who have conducted previous reintroduction attempts.

Justification for inclusion

The Fen Orchid is one of the most endangered wildflowers in Europe. Formerly known from at least 30 fenland sites in eastern England in the 1890s, only three native populations of Fen Orchid are known to survive in the Norfolk Broads.

After a decade of research and partnership work, the orchid has been re-discovered at former sites in the Broads, and the total population has estimated to have risen to over 15,000 plants through proper management.

The orchid has also been reintroduced to its former sites in Suffolk, and the signs are encouraging that it will become established in some of its old homes.

Habitat grouping

Lowland fens, floodplain wet grasslands and associated habitats.



Fen Orchid



Did you know?

The first ever local flora (a book about local plants) was about Cambridge and appeared in 1660.



8

Local habitat map

Local habitat map

The local habitat map is available online in the form of a set of interactive maps.

These maps allow users to understand the priorities in their locality and inform the most effective action for nature recovery across Cambridgeshire and Peterborough.



<https://experience.arcgis.com/experience/a1963a7dc51c4c259a8d19832b63f461>

Habitat priorities / Methodology

For the habitat priorities, we created a map of areas that are already important for biodiversity. This map is a collation of Sites of Special Scientific Interest (SSSIs) and Local Nature Reserves.

We then created a map of areas where there are further opportunities for nature recovery.

To do this we:

- 1 Asked people across Cambridgeshire and Peterborough what habitats and species were important to them.
- 2 Created a map of existing habitats across Cambridgeshire and Peterborough. These habitats include areas of woodland, wetland, grassland, farmland, the urban landscape, rivers, streams and drains
- 3 Used computer modelling and a wide range of datasets to identify how and where habitats could be created or enhanced
- 4 Ensured existing land uses were considered, including the presence of buildings or historic features and connectivity of habitats across the landscape etc
- 5 Prioritised areas of land where we identified opportunities to create a 'buffer' around existing nature sites or where they could provide habitat 'stepping stones' between existing habitats including nationally and locally designated sites
- 6 Mapped ecosystem service benefits such as where land has the potential to improve air or water quality, provide carbon storage or manage flood risk
- 7 Further prioritised areas where both habitat and ecosystem services benefits could be demonstrated
- 8 Developed a single map of the best opportunities for each habitat priority and working with local experts and farmer representative organisations to sense-check these decisions. As part of this we incorporated large scale priority natural landscapes which have previously been identified by work undertaken by Natural Cambridgeshire

For the species priorities we worked with local ecologists, experts and other stakeholder input to create a list of priority species based on the threat posed to them, their significance to the region, and the deliverability of any intervention.

Actions that would help contribute to their recovery or enhancement were then developed.

Next steps

Nature recovery requires positive action to be taken by all of us. The scale of the challenge, the numbers of actions and the area they cover means it will not be possible to move forward on all fronts at once.

While some actions in some areas are already well advanced, and others have recently started or are being planned, many of the actions by their nature are long-term, and beneficial outcomes will not be apparent in the short-term.

To be effective the Local Nature Recovery Strategy will need a **clear and prioritised delivery plan**, setting out which actions are prioritised, the locations where they are being prioritised, and who is delivering them.

For some of the longer-term actions, manageable milestones will need to be set, perhaps with SMART short-term objectives and actions.

The **delivery plan** will need to cross-reference with other sectoral plans and be regularly updated as and when new actions commence, or actions are completed. It should be capable of being presented by location (action is always local to somewhere) or by habitat or species categories.

Monitoring & Measurement

We also need to be able to measure and report on success. Monitoring must include measurement of real biological and ecological outcomes. The following elements will need to be measured:

- **Habitat Quality:** Changes in the ecological condition of high-value nature sites (SSSIs and

Local Wildlife Sites). (This can be measured relatively quickly and updated annually).

- **Habitat Area:** Changes in area and % land cover of high-value habitats. (This is best measured at set periods e.g. every 10-20 years, though with a good baseline, incremental changes could be recorded from multiple data sources).
- **Habitat Connectivity:** Increases in habitat connectivity and conversely decreases in habitat fragmentation. (This is likely to require habitat modelling methods at a county-scale and would be repeated on a similar timescale to monitoring of the extent of high-value habitats e.g. every 10-20 years).
- **Species:** Species can be monitored in terms of numbers of species found in our area, the populations of individual species and overall bio-abundance. (Species are the most challenging measure, but in the short-term it may be possible to use national datasets to get local data for some species groups, e.g. breeding birds and butterflies. Bespoke surveys and monitoring could also be undertaken for those LNRS priority species where action is being taken forward).

However, at the scale of a whole county some of the above measures will only be updated every 10 or more years. Measuring progress will therefore also need to record actions taking place in specific locations, which could demonstrate increased efforts and impact over time, pending results from county-wide programmes for measuring changes in habitats and species.



Acknowledgements

We thank the following organisations and individuals who have helped develop this draft LNRS for consultation through attending workshops and/or providing information.

Anglian Water

Cam and Ely Ouse Catchment Partnership

Cambridge City Council

Cambridge Past, Present and Future

Cambridge Water

Cambridgeshire and Peterborough
Environmental Records Centre

Cambridgeshire County Council

Cambridgeshire Geological Society

Centre for Landscape Regeneration, Cambridge
University

Country Land and Business Association

Diocese of Ely

East Cambridge Farmers

East Cambridgeshire District Council

Eco Church

Ely Nature-Friendly Farming Zone

Environment Agency

Farming and Wildlife Advisory Group (FWAG)

Fenland District Council

Fenland SOIL

Forestry Commission

GeoPeterborough

Great Ouse Rivers Trust

Greater Cambridge Planning

Huntingdonshire District Council

John Clare Countryside

Keystone Marketing

Land Use Consultants

Middle Level Commissioners, Internal Drainage
Board

National Farmers Union

National Trust

Natural Cambridgeshire

Natural Capital Solutions

Natural England

Nene Park Trust

Ouse Valley Farm Cluster

Peterborough City Council

Royal Society for the Protection of Birds (RSPB)

Respondents to all public surveys and
questionnaires

South Cambridgeshire District Council

Thornley Farm Cluster

Urban and Civic

Water Resources East

West Camb 100s

Wildfowl & Wetlands Trust

Wildlife Trust for Bedfordshire, Cambridgeshire
and Northamptonshire

Woodland Trust

Section five

Page 1

Worts Meadow Local Nature Reserve
© Paul England Photography

Page 2

Great Raveley Drain © Pamela Abbott
Hedgehog © Tom Marshall (WildNet)
Water buffalo at Chippenham Fen
© Pamela Abbott
Small Bloody-nose Beetle © Paul Rule
Sedge Fen sunset © Sally Fisher

Page 3

Wicken Fen sunrise
© National Trust, Rob Coleman

Page 4

Accessibility with tramper at Wandlebury
© Pamela Abbott
Vestal Cuckoo Bumblebee (Cambourne)
© Brian Eversham
Urban landscape, Cambridge © Pamela Abbott
Background photo Chippenham Fen
© Pamela Abbott
Station Wood in Isleham
© Paul England Photography

Page 5

Barn Owl © Natural Cambridgeshire
Background photo Chippenham Fen
© Pamela Abbott

Page 6

Connington & Holme Wood
© Paul England Photography

Page 7

Commissioners' Drain sunset © Pamela Abbott

Page 8

Great Fen ditch with reeds and trees
© Pamela Abbott

Page 9

Ouse Fen Drone shot © Jeff Kew

Page 10

Felmersham Gravel Pits © Noa Leach
Fen Drayton Lakes © Hilary Tandy

Page 11

Background image © Paul England Photography
Wicken Fen © Pamela Abbott
John Clare Countryside © Pamela Abbott

Nene Park © Rob Harradine

Page 12

Ouse washes © Pamela Abbott
Fulbourn Fen © Pamela Abbott
West Cambridgeshire Hundreds © Pamela Abbott

Page 13

Tansy beetle, Woodwalton Fen © Pamela Abbott
Washingley © Rob Harradine

Page 14

Background image Worts Meadow Local Nature
Reserve © Paul England Photography

Page 15

Fleam Dyke © Pamela Abbott
Nene Wetlands © Robert Enderby

Page 16

Crested Cow-wheat, Claylands © Brian Eversham
Babraham hedge © CWS Surveys
Green-winged Orchid © Brian Eversham

Page 17

Pasque Flower, Devil's Dyke © Paul Rule
Chalk Hill Blue male, Devil's Dyke © Paul Rule

Page 18

City of Ely drone shot © Paul England Photography

Section six

Page 19

Dark Green fritillary © Paul Rule

Page 20

Background image Forest landscape at Hill Farm
© Paul England Photography
Skylark © David Tipling 2020Vision

Water Vole in the reeds, Ring end
© For photography accreditation please contact
Cambridgeshire County Council

Page 21

Knapwell Wood © Pamela Abbott

Page 22

Ninewells Local Nature Reserve © Pamela Abbott
Overhall Grove Nature Reserve © Pamela Abbott

Page 23

Hatley Elm © Brian Eversham
Wet woodland © Pamela Abbott
Orchard blossom © Pamela Abbott

Page 24

Bluebells in Brampton Wood © Sophie Baker

Oxlip in Waresley, Gransden Woods
© Robert Enderby

Bluebells © Brian Eversham

Purple Emperor Butterfly © Mike Addis

Hazel Dormouse © Gwen Hitchcock

Page 25

Bluebells in Brampton Wood © Sophie Baker

Purple Emperor © Mike Addis

Page 25

Green-winged Orchid, Upwood Meadows © Brian Eversham

Pasque Flower © Pamela Abbott

Page 26

Knapweed Broomrape © Henry Stanier

Waxcap © Brian Eversham

Glow-worm © Brian Eversham

Page 27

Darlow Farm © Holly Wilkinson

Page 28

Woodwalton Fen © Pamela Abbott

Spined Loach © Brian Eversham

Paludiculture Beds Great Fen © Holly Wilkinson

Page 29

Gordons Mere © Thomas Sisman

Fenland drain ditches © Sofi Lloyd

Page 30

Dernford Reservoir, Stapleford © Pamela Abbott

Ground Beetle © Brian Eversham

Fen Violet © Brian Eversham

Yellow Iris © Brian Eversham

Drinker moth caterpillar © Brian Eversham

Marsh Fragrant Orchid © Brian Eversham

Page 31

Wicken Fen © Pamela Abbott

Page 32

The Ouse Washes © Paul England Photography

Page 33

River Nene - old course © Rob Harradine

Spined Loach © Brian Eversham

Opposite-leaved Pondweed © Brian Eversham

River Snail © Brian Eversham

Page 34

River Granta drone shot

Page 35

Holme Fen (Autumn) © Sarah Lambert

Skylark © David Tipling 2020Vision

Common Lizard © Guy Pilkington

Page 36

Play area, Alconbury green corridor, Alconbury Weald © Urban&Civic

Pond dipping with Mike Dilger © Urban&Civic

Grazing cows in front of the Ely Cathedral
© Jakub Pabis

Page 37

Urban landscape, Northstowe © Pamela Abbott

Urban landscape, Alconbury Weald © Urban&Civic

Page 38

Hedgehog © Tom Marshall

Brimstone butterfly © Pamela Abbott

Swallow © Chris Gomersall 2020Vision

Bee Orchid Brian Eversham

Cambridge © Paul England Photography

Page 39

Farmed landscape, Ely © NFU

Page 40

Upwood Meadows © Brian Eversham

Belted Galloway conservation grazing,
Wandlebury © Pamela Abbott

Page 41

Habitat mosaic, Cambourne © Pamela Abbott

Farmed landscape hedges, Fulbourn
© Pamela Abbott

Harvest Mouse © Colin Bradshaw

Farmed landscape, Stapleford © Pamela Abbott

Corn Marigold © Brian Eversham

Broad-leaved spurge © Brian Eversham

Page 42

Barnack Hills and Holes © Pamela Abbott

Hayley Wood, West Cambridgeshire Hundreds
© Pamela Abbott

Ouse valley © Pamela Abbott

Page 43

Holme Fen, Caldecote Dyke © Rob Harradine

Holme Fen, Stilton Dyke © Rob Harradine

Gog Magog Downs © Pamela Abbott

Page 44

Southern and Early Marsh Orchids at Fulbourn Fen
© Pamela Abbott

Bluebell and Stichwort at Brampton Wood ©
Pamela Abbott

Page 45

Chippenham Fen National Nature Reserve ©
Pamela Abbott

Grassland management on Devil's Dyke
© Jenny Payne

Section seven

Page 46

Farmed landscape, arable field margin
© Pamela Abbott

Page 47

Black-tailed Godwit © Neil St John

Stone Curlew © RSPB, Andy Hay

Page 48

Swift © David Tipling 2020Vision

Turtle Dove © For photography accreditation
please contact Cambridgeshire County Council

Page 49

Fairy Shrimp - image to follow

White-clawed Crayfish © Linda Pitkin 2020Vision

Page 50

Autumn Lady's-tresses © Brian Eversham

Bastard-toadflax © Brian Eversham

Broad-leaved Cudweed © Emily Swan

Page 51

Cambridge Milk Parsley © Simon Stirrup

Common Fragrant Orchid © Brian Eversham

Common Butterwort © Brian Eversham

Page 52

Dodder © Paul Rule

Common Cottongrass © Brian Eversham

Crested Cow-wheat © Brian Eversham

Page 53

Dandelion © Glynis Pierson

Fen Wood-rush, Woodwalton Fen © Sarah Lambert

Page 54

Fibrous Tussock-Sedge © Sarah Lambert

Flat-sedge © Malcolm Storey, 2008
www.bioimages.org.uk

Page 55

Fly Orchid © Brian Eversham

Frog Orchid © Pat Doody

Great Pignut © Alex Prendergast

Page 56

Goldilocks Buttercup © Brian Eversham

Page 57

Fen form of Heath Dog-violet © Brian Eversham

Page 58

Maiden Pink (with Hoverfly, Sphaerophoria
scripta) © Rob Dyke

Marsh Fragrant-Orchid © Brian Eversham

Page 59

Marsh Helleborine © Brian Eversham

Marsh Lousewort © Brian Eversham

Page 60

Milk Parsley © Simon Stirrup

Moon Carrot © Brian Eversham

Page 61

Mountain Everlasting © Brian Eversham

Pasque Flower © Paul Rule

Page 62

Rare Spring-sedge © Sarah Lambert

Red-tipped Cudweed © J.Masson

Page 63

Slender Hare's-ear © Sarah Lambert

Smooth Rupturewort © J.Masson

Spanish Catchfly © Sarah Lambert

Page 64

Spotted Cat's-ear © John Secker

Violet Helleborine © Brian Eversham

Page 65

Black Hairstreak © Brian Eversham

Caddisfly Trichoptera © Chris Lawrence

Page 66

Eyed Longhorn Beetle © Brian Eversham

Desmoulin's Whorl Snail © Simon Stirrup

Page 67

Ground Beetle © Brian Eversham

Large-mouthed Valve Snail, Woodwalton Fen
© Sarah Lambert

Page 68

Marsh Carpet Moth © Simon Stirrup

Mud Beetle - image unavailable

Page 69

Pardosa Paludicola Spider - image to follow

Tansy Beetle © Brian Eversham

Page 70

Taylor's Spire Snail - For photography accreditation please contact Cambridgeshire County Council

Witham Orb Mussel - For photography accreditation please contact Cambridgeshire County Council

Page 71

Zora Armillata Spider © Jorge Almeida

Hazel Dormouse © For photography accreditation please contact Cambridgeshire County Council

Page 72

Adder © Trevor Baker

Lichen (*Bellicidia incompta*) © Mark Powell

Page 73

Hayley Elm © Brian Eversham

Page 74

Bearded Stonewort, Orton Pit © Sarah Lambert

Dwarf Stonewort © Chris Carter

Page 75

Tassel Stonewort © Chris Carter

Habitat mosaic, Nene Park © Pamela Abbott

Section eight

Page 76

Corncrake © RSPB

Fen Ragwort © Brian Eversham

Page 77

Fen Violet © Brian Eversham

Large Marsh Grasshopper © Brian Eversham

Page 78

Water Germander © Simon Stirrup

Pool Frog © Amphibian and Reptile Conservation

Page 79

Large Copper butterfly © Natural England

Swallowtail butterfly © Stefan Johansson

Page 80

Fen Orchid © Malcolm Storey, 2003
www.bioimages.org.uk

Ouse Fen © Pamela Abbott

Section nine

Page 81

Snowdrops © National Trust, James Beck

Page 83

Sawtry Fen © Rob Harradine

Page 89

Photo competition winner, Ely sunrise
© Chris Barton



Cambridgeshire and Peterborough
LOCAL NATURE RECOVERY STRATEGY

Enquiries email localnaturerecoverystrategy@cambridgeshire.gov.uk
or visit NaturalCambridgeshire.org.uk/lnrs